

Utility of Silver Nitrate Test for Post-Mortem Diagnosis of Aluminum Phosphide Poisoning

Ali Banagozar Mohammadi¹, Maryam Zaare Nahandi*¹, Hassan Solhi²

Received: 06.12.2012

Accepted: 23.12.2012

ABSTRACT

Background: Intoxication with aluminum phosphide has been prevalent in Iran recently. Lab confirmation of the poisoning is needed in clinical and post-mortem settings. Present study was conducted to determine efficiency and validity of silver nitrate test for detection of aluminum phosphide in samples.

Methods: Information about dead patients with history of aluminum phosphide poisoning referred for medicolegal autopsy was collected during 18 months. Their relationships with result of silver nitrate test was assessed statistically.

Results: 24 patients were studied. Silver nitrate test on samples of intra-abdominal organs was positive in 75% of autopsies, while the test on gastric content was positive in 50%. Relationship between number of ingested tablets and silver nitrate test was statistically significant (p value=0.017).

Conclusion: Results of present study showed that silver nitrate test can be used as a preliminary method for evaluation of patients or dead bodies suspected to be intoxicated with ALP. However, even negative result of silver nitrate test would not preclude ALP poisoning and more definitive quantitative tests such as HS-GC should be ordered, if feasible.

Keywords: Aluminum Phosphide, Diagnosis, Intoxication, Silver Nitrate, Test.

IJT 2013; 804-807

INTRODUCTION

Intoxication with pesticides is a worldwide health problem, particularly in developing countries. It causes more than 300,000 deaths annually, mainly after intoxication with organophosphates, organochlorines and aluminum Phosphide (1). Aluminum phosphide (ALP) is an effective and economical pesticide (2). It is an inorganic solid substance with dark grey or yellow crystals. Its melting point is more than 1000°C. It is heavier than water. It reacts with water, produces phosphine, a highly toxic gas (3-5). Phosphine is an appropriate protective factor according to international biological protocols. It is widely used for protection of crops, grains,

and commodities because of its effectiveness, wide accessibility, and low cost (6). Phosphine, also named hydrogen Phosphide and celphos, is a colorless, flammable and explosive gas with rotten fish or garlic odor. Its molecular weight is 34 g/mol with melting point -133.5 °C and boiling point -87.5 °C (7,8).

Recently ALP in form of 3 grams tablet contains 1680 mg aluminum phosphide and 1320 mg ammonium carbonate is widely used for suicidal attempts by Iranian youths (9-14). It poses a new health dilemma. Ingestion of only one-fourth tablet is lethal. There are reports of death after ingestion of 150-500 mg ALP (15-17). Its LD50 is estimated to be

1. Department of Internal Medicine, Tabriz University of Medical Sciences, Tabriz, Iran.

2. Department of Clinical Toxicology and Forensic Medicine, Arak University of Medical Sciences, Arak, Iran.

*Corresponding Author E-Mail: dr_mzaare@yahoo.com

20 mg/kg (18). ALP poisoning creates multi-organ failure and variable clinical presentations (19).

Routinely chemical tests are used for confirmation of the intoxication, if patient's history or clinical presentation is warranted. Qualitative tests are used for verification of presence of phosphine in blood, breath, gastric contents, and other tissues and quantitative tests are used for determination of severity or prognosis of poisoning (8,16). There are qualitative (such as colorimetry), quantitative (for example GC, ETAAs, and ICP-MS) and semi-quantitative (potentiometry) methods for detection of ALP in the samples (2,8,19,20). Colorimetric methods such as silver nitrate test can be performed on gastric aspirate, exhaled breath, and non-biologic samples. If a paper immersed with silver nitrate is kept in contact with a questioned sample contained ALP or phosphine, it would turn black (2,21,22).

Silver nitrate test is performed in autopsy ward of Iranian legal medicine organization for cases suspected to die after ALP intoxication. But how much can actually rely on results of the test? Present study was conducted to determine efficiency and validity of the test.

MATERIALS AND METHODS

During 18 months, from Sep.2009 to Jan. 2011 all dead bodies that referred to autopsy ward of Iranian legal medicine organization for medicolegal autopsy with history of ALP poisoning were included in the study. Information about chief complaint, vital signs, clinical presentation, dose and route of ingestion, demographic characteristics, lab findings, duration of hospital admission were extracted from hospital records of the patients. Findings in autopsy of the patients include macroscopic and histopathologic findings as well as result of silver nitrate test on gastric content and intra-abdominal tissues, and GC and TLC tests for detection of other poisons and drugs in the biologic samples were recorded too. The collected

data were analyzed using SPSS software version 16.0.

RESULTS

24 patients with above mentioned criteria were included in the study include 8 (33.3%) males and 16 females. 5 patients had history of previous psychiatric problem; while 4 had previous suicidal attempts and one was drug abuser. All but one had been intoxicated after suicidal attempts. Mean number of ingested tablet was 1.29 ± 0.54 (1-3 tablets). Mean age of the patients was 24.12 ± 10.41 (14-66 years old). The most common symptom after arrival to the hospital was nausea and vomiting.

The autopsies had been performed within 48 hours after death. Body decomposition had not occurred at the time of autopsy. Silver nitrate test on samples of intra-abdominal organs was positive in 75% of autopsies, while the test on gastric content was positive in 50%. Mean number of ingested tablet was 1.43 ± 0.62 in autopsies with positive silver nitrate test on samples of intra-abdominal organs. Patients with negative silver nitrate test on samples of intra-abdominal organs had ingested only one tablet of ALP.

Relationship between number of ingested tablets and silver nitrate test was statistically significant (p value=0.017). There were no statistically significant relationships between other evaluated parameters with result of silver nitrate test.

DISCUSSION

Results of present study showed that silver nitrate test can be used as a preliminary method for evaluation of patients or dead bodies suspected to be intoxicated with ALP.

Various studies have suggested that if fresh tablets of ALP have been responsible for the intoxication, sensitivity of silver nitrate test reaches to 100% for samples of gastric contents and 50% for samples of exhaled breaths (23,24). Efficiency of the test is estimated 90% (8).

However we reach the sensitivity 75 and 50 percent for post-mortem samples from intra-abdominal organs and gastric content respectively. It should not be forgotten that negative silver nitrate test do not rule out ALP poisoning, particularly in post-mortem settings.

CONCLUSION

Silver nitrate test can be used as a preliminary test for assessment of likelihood of ALP poisoning. Nevertheless, if patient's history, clinical presentation, and autopsy findings were highly suggestive of ALP poisoning, even negative result of silver nitrate test would not preclude ALP poisoning and more definitive quantitative tests such as HS-GC should be ordered, if feasible (2,20,25,26).

ACKNOWLEDGMENTS

The authors extend their thanks from Maryam Akhgari (head officer of toxicology laboratory of Iranian legal medicine organization) for her contribution in performing silver nitrate test in the study.

REFERENCES

- Goel A, Aggarwal P. Pesticide poisoning. National Medical Journal of India. 2007;20(4):182-91.
- Galichet LY, Moffat A, Osselton M, Widdop B. Clarke's analysis of drugs and poisons. Pharmaceutical Press, London, England; 2004.
- Anand R, Binukumar B, Gill KD. Aluminum phosphide poisoning: an unsolved riddle. Journal of Applied Toxicology. 2011;31(6):499-505.
- Inchem.org [homepage on the internet]. Chemical Safety Information from Intergovernmental Organizations, Aluminum phosphide. available from : <http://www.inchem.org/documents/icsc/icsc/eics0472.htm>
- Exttoxnet.orst.edu [homepage on the internet]. Pesticide Information Profiles. available from: <http://exttoxnet.orst.edu/pips/alumphos.htm>
- Valmas N, Ebert PR. Comparative toxicity of fumigants and a phosphine synergist using a novel containment chamber for the safe generation of concentrated phosphine gas. PloS one. 2006;1(1):e130.
- Phosphine; Determination of Noncancer Chronic Reference Exposure Levels; Office of Environmental Health Hazard Assessment. available from : http://oehha.ca.gov/air/chronic_rels/pdf/7803512.pdf
- Inchem.org [homepage on the internet]. Chemical Safety Information from Intergovernmental Organizations, Phosphine. available from : <http://www.inchem.org/documents/icsc/icsc/eics0472.htm>
- Mehrpour O, Alfred S, Shadnia S, Keyler D, Soltaninejad K, Chalaki N, et al. Hyperglycemia in acute aluminum phosphide poisoning as a potential prognostic factor. Human & experimental toxicology. 2008;27(7):591-5.
- Shadnia S, Rahimi M, Pajoumand A, Rasouli MH, Abdollahi M. Successful treatment of acute aluminium phosphide poisoning: possible benefit of coconut oil. Human & experimental toxicology. 2005;24(4):215-8.
- Saleki S, Ardalan FA, Javidan-Nejad A. Liver histopathology of fatal phosphine poisoning. Forensic science international. 2007;166(2):190-3.
- Mehrpour O, Dolati M, Soltaninejad K, Shadnia S, Nazparvar B. Evaluation of histopathological changes in fatal aluminum phosphide poisoning. cortex. 2008;26:57.8.
- Soltaninejad K, Faryadi M, Sardari F. Acute pesticide poisoning related deaths in Tehran during the period 2003–2004. Journal of forensic and legal medicine. 2007;14(6):352-4.
- Mostafazadeh B, Pajoumand A, Farzaneh E, Aghabiklooei A, Rasouli MR. Blood levels of methemoglobin in patients with aluminum phosphide poisoning and its correlation with patient's outcome. Journal of Medical Toxicology. 2011;7(1):40-3.
- Gupta S, Ahlawat SK. Aluminum phosphide poisoning-a review. Clinical Toxicology. 1995;33(1):19-24.
- WHO. Aluminium phosphide - draft revised Poisons Information Monograph for peer review. . available from:

- http://www.who.int/ipcs/poisons/pim_aluminium_phosphide.pdf
17. Wahab A, Zaheer M, Wahab S, Khan R. Acute aluminium phosphide poisoning: an update. *Hong Kong J Emerg Med.* 2008;15(3):152-5.
 18. Bajwa S, Bajwa Kaur S, Kaur J, Singh K, Panda A. Management of celphos poisoning with a novel intervention: A ray of hope in the darkest of clouds. *Anesthesia: Essays and Researches.* 2010;4(1):20-4.
 19. Bumrah GS, Krishan K, Kanchan T, Sharma M, Sodhi GS. Phosphide poisoning: A review of literature. *Forensic science international.* 2012;214(1):1-6.
 20. Musshoff F, Preuss J, Lignitz E, Madea B. A gas chromatographic analysis of phosphine in biological material in a case of suicide. *Forensic science international.* 2008;177(2):e35-e8.
 21. Sudakin D. Occupational exposure to aluminium phosphide and phosphine gas? A suspected case report and review of the literature. *Human & experimental toxicology.* 2005;24(1):27-33.
 22. Tripathi S, Gautam C, Sharma P. Clinical pharmacology of aluminium phosphide poisoning. *Indian Journal of Pharmacology.* 1992;24(3):134-7.
 23. Chugh S, Ram S, Chugh K, Malhotra K. Spot diagnosis of aluminium phosphide ingestion: an application of a simple test. *The Journal of the Association of Physicians of India.* 1989;37(3):219-20.
 24. Mital H, Mehrotra T, Dwivedi K, Gera M. A study of aluminium phosphide poisoning with special reference to its spot diagnosis by silver nitrate test. *Journal of Association of Physicians of India.* 1992;40:473-4.
 25. Gurjar M, Baronia AK, Azim A, Sharma K. Managing aluminum phosphide poisonings. *Journal of Emergencies, Trauma and Shock.* 2011;4(3):378-84.
 26. Chan L, Crowley R, Delliou D, Geyer R. Phosphine analysis in post mortem specimens following ingestion of aluminium phosphide. *Journal of analytical toxicology.* 1983;7(4):165-7.