

Original Article**Acute Toxicity of a Heavy Metal Cadmium to an Anuran, the Indian Skipper Frog *Rana cyanophlyctis***Ajai Kumar Srivastav*¹, Shilpi Srivastav¹, Nobuo Suzuki²

Received: 29.03.2016

Accepted: 16.04.2016

ABSTRACT

Background: There has been increasing awareness throughout the world regarding the remarkable decrease in amphibian population. For such amphibian population decline several causes have been given. Cadmium, a heavy metal is released both from natural sources (leaching of cadmium rich soils) and anthropogenic activities to the aquatic and terrestrial environments. This study evaluated the toxicity of heavy metal cadmium to Indian skipper frog *Rana cyanophlyctis*.

Methods: For the determination of LC₅₀ values for cadmium, four-day static renewal acute toxicity test was used. Five replicates each containing ten frogs were subjected to each concentration of cadmium chloride (15, 20, 25, 30, 35, 40, 45 and 50 mg/L). At different exposure periods (24, 48, 72 and 96 h), the mortality of the frog was subjected to Probit analysis with the POLO-PC software (LeOra Software) to calculate the LC₅₀ and 95% confidence level.

Results: The LC₅₀ values of cadmium chloride for the frog *R. cyanophlyctis* at 24, 48, 72, and 96 h are 32.586, 29.994, 27.219 and 23.048 mg/L, respectively. The results have been discussed with the toxicity reported for other aquatic vertebrate --fish.

Conclusion: Cadmium caused mortality to the frog and this could be one of the reasons for population decline of frogs which inhabit water contaminated with heavy metals.

Keywords: Amphibia, Cadmium, Heavy Metal, LC₅₀, Toxicity.

IJT 2016 (5): 39-43

INTRODUCTION

The heavy metals are discharged into water resources through industries such as steel and mining, textile dyes, paints and varnishes, fertilizers, feed additives and ceramics [1]. "The global dispersion of heavy metals and their extensive use in modern society pose an enormous challenge to organisms including humans" [2]. In the aquatic and terrestrial environments the release of heavy metal cadmium is through leaching of cadmium rich soils, mining, smelting, electroplating, manufacturing of batteries etc. [3]. "Atmospheric deposition of airborne cadmium, and the application of cadmium-containing fertilizers and sewage sludge on farm land may lead to contamination of soils and increased cadmium uptake by crops and vegetables consumed by human beings" [4]. Cadmium has been considered more toxic (toxic at levels one tenth) that of lead, mercury, aluminium or nickel [5] and ranked the 7th toxicant in the Priority List

of Hazardous Substances of the Agency for Toxic Substances and Disease Registry [6, 7]. This element has a long-biological half-life in humans and it gets accumulated in vital organs – especially in liver and kidney throughout their lives [8, 9].

There has been increasing awareness throughout the world regarding the remarkable decrease in amphibian population [10, 11]. A comprehensive global assessment of amphibian species has been performed by The International Union for Conservation of Nature (IUCN)[12]. Almost 32.5% were listed as vulnerable, endangered, or critically endangered. 7.4% species were listed as Critically Endangered, and about 43% were experiencing some form of population decline [13]. For such amphibian population decline several causes have been given which are habitat loss [12]; UV-B radiation [14,15]; infectious disease [16, 17]; contaminants [18-20]; non-native predators [21]; a combination

1. Department of Zoology, DDU Gorakhpur University, Gorakhpur 273 009, India.

2. PhD. Noto Marine Laboratory, Institute of Nature and Environmental Technology, Kanazawa University, Ogi, Noto-cho, Ishikawa 927-0553, Japan.

*Corresponding Author: E-mail: ajaisrivastav@hotmail.com

of factors [22, 23] and pesticide's effect [19, 20, 24]. There is a wide variation in tolerance levels among amphibians even between closely related species [25]. Conclusions drawn from studies on only a few species cannot reveal the entire effects of any harmful chemicals to amphibians in general [26].

Hence, in this study we have assessed the LC₅₀ value of heavy metal cadmium on an anuran species *Rana cyanophlyctis*. This species has been chosen for the experiments as this frog is highly aquatic and remains permanently resident in different types of habitats with pooled water.

MATERIALS AND METHODS

For experiments laboratory bred *R. cyanophlyctis* (both sexes, body wt. 14.74 ± 0.55 g) were selected. Mean weights of the frogs used in the experiment showed no significant difference. Frogs were kept in all glass aquaria containing dechlorinated tap water and acclimatized to the laboratory conditions (under natural photoperiod 11.58-12.38 and temperature 27.2 ± 1.4 °C) for at least two weeks. The physicochemical characteristics of the tap water were pH 7.20 ± 0.1 ; dissolved oxygen 7.95 ± 0.25 mg/L and hardness as CaCO₃ 167.06 ± 5.61 mg/L. Frogs were fed daily with live insects, 2-3 times per day. Water was renewed daily after cleaning the fecal matter. Feeding was stopped 24 h before and during the experimental period to avoid the excretory substances to influence the toxicity test solutions.

All the experimental protocols were approved by the Ethical Committee of Department of Zoology, DDU Gorakhpur University.

For the determination of LC₅₀ values for cadmium, four-day static renewal acute toxicity test [27] was used. Five replicates each containing ten frogs (kept in glass aquarium containing 30 L of the test solution) were subjected to each concentration of cadmium chloride (15, 20, 25, 30, 35, 40, 45 and 50 mg/L). Cadmium chloride was firstly dissolved in distilled water and then desired volume of the solution was mixed in tap water to obtain the above mentioned toxicant concentrations. For each toxicant, a control group with five replicates (each containing 10 frogs) kept in 30 L tap water was also run. The solutions of all the aquaria (control and experimental) were renewed daily. Precautions were taken to remove the dead frog immediately because dead animals

deplete dissolved oxygen which greatly affects toxicity data. At different exposure periods (24, 48, 72 and 96 h), the mortality of the frog was subjected to Probit analysis with the POLO-PC software (LeOra Software) to calculate the LC₅₀ and 95% confidence level.

RESULTS

The per cent mortality of *R. cyanophlyctis* after exposure to various concentrations of cadmium chloride for 24, 48, 72 and 96 h is shown in Figures 1-4. The LC₅₀ values (Table 1) for cadmium chloride at 24, 48, 72 and 96 h are 32.586, 29.994, 27.219 and 23.048 mg/L, respectively. Table 1 also depicts the slope functions and upper and lower confidence limits for *R. cyanophlyctis*.




Figure 1. Per cent mortality of the frog *Rana cyanophlyctis* after 24 h exposure to different concentrations of cadmium chloride.




Figure 2. Per cent mortality of the frog *Rana cyanophlyctis* after 48 h exposure to different concentrations of cadmium chloride.




Figure 3. Per cent mortality of the frog *Rana cyanophlyctis* after 72 h exposure to different concentrations of cadmium chloride.




Figure 4. Per cent mortality of the frog *Rana cyanophlyctis* after 96 h exposure to different concentrations of cadmium chloride.

Table 1. LC₅₀ value, slope function and confidence limits for short-term exposure of cadmium chloride at different intervals for the frog *R. cyanophlyctis*.

| Exposure Periods | Effective dose (mg/L) | Limits(mg/L)* | | Slope Function | 't' ratio | Heterogeneity |
|------------------|--------------------------|---------------|--------|----------------|-----------|---------------|
| | | LCL | UCL | | | |
| 24 h | LC ₁₀ =22.858 | 19.731 | 25.163 | 8.322 | 12.652 | 1.687 |
| | LC ₅₀ =32.586 | 30.415 | 34.800 | ± | | |
| | LC ₉₀ =46.454 | 42.502 | 53.089 | 0.658 | | |
| 48 h | LC ₁₀ =21.121 | 17.646 | 23.598 | 8.413 | 13.101 | 2.307 |
| | LC ₅₀ =29.994 | 27.555 | 32.407 | ± | | |
| | LC ₉₀ =42.594 | 38.644 | 49.553 | 0.642 | | |
| 72 h | LC ₁₀ =18.448 | 15.782 | 20.504 | 7.586 | 13.258 | 1.532 |
| | LC ₅₀ =27.219 | 25.225 | 29.148 | ± | | |
| | LC ₉₀ =40.161 | 36.823 | 45.371 | 0.572 | | |
| 96 h | LC ₁₀ =16.153 | 13.485 | 18.115 | 8.301 | 12.571 | 1.863 |
| | LC ₅₀ =23.048 | 21.085 | 24.895 | ± | | |
| | LC ₉₀ =32.886 | 30.010 | 37.587 | 0.660 | | |

*The upper and lower confidence limits for LC₅₀ values calculated at 0.05 levels.

DISCUSSION

In the present study the LC₅₀ values for cadmium chloride for adult *R. cyanophlyctis* were 32.58, 29.99, 27.21 and 23.04 at 24, 48, 72 and 96 h, respectively. The percentage mortality of the frogs increased as the concentration of cadmium increased. This increase was also time dependent. This is consistent with the reports of Ezemonye and Enuneku [28] who have also noticed increased mortality of tadpoles of *Bufo maculatus* and *Ptychudena bibroni* with increased lead concentration.

“The mean LC50 value in *Bufo. maculates* for 24 h were not determined due to low mortality. However, LC50 values for 48, 72 and 96 hours were 17.74, 13.39 and 9.97mg/l respectively”

[29]. Woodal et al. [30] have reported 90 h LC₅₀ of cadmium for *Xenopus laevis* larvae between 80 and 100 mg/L. 48 h LC₅₀ value for *Xenopus laevis* larvae has been found as 11.7 mg/L [31] and 7.36 mg/L [32]. 96 h LC₅₀ value for cadmium for *Bufo arenarum* larvae has been reported as 2.19- 6.77 mg/L [33]. Sparling and Lowe [18] have reported LC₅₀ value of cadmium for *Rana clamitans* as 1.9 mg/L. Grillitch and Chovanec [34] recorded LC₅₀ value of cadmium as 0.45 mg/L for larvae of *Rana ridibunda*. For *Rana ridibunda* tadpoles the 96 h LC₅₀ has been reported as 71.8 mg/L [35]. 96 h LC₅₀ value for cadmium for larvae of *Rana ridibunda* has been reported as 51.2 mg/L [11].

The toxicity of cadmium for teleosts has been studied extensively. The 96 h LC₅₀ value of cadmium for various teleosts has been reported by

several investigators as – 46.8 mg/L for *Carassius auratus* [36], 0.63 (in soft water) and 73.5 mg/L (in hard water) for *Pimephalus promelas* [37], 3.15 mg/L for *Notemigonus crysoleucas* [38], 6.72 mg/L for *Barilius vagra* [39], 89.5 mg/L for *Labeo rohita* [40] and 360 mg/L for *Heteropneustes fossilis* [41].

The toxicity of cadmium to various amphibians has been studied mostly in tadpole/larval stages. There exists studies on LC₅₀ value of cadmium for amphibians which reported the values as 50.0 mg/kg (for 48 h in males of toad) [42] and as 9.90 mg/L (for 96 h for *Bufo maculatus*) [29]. In the present study the 96 h LC₅₀ for cadmium for adult *R. cyanophlyctis* has been found as 23.04 mg/L. This indicates that *R. cyanophlyctis* is more resistant/tolerant to cadmium as compared to *Bufo maculatus*. "Tolerance is an important mechanism by which an organism reacts to an adverse environment. Mechanisms that might be responsible for tolerance include decreased uptake, metal speciation, increased excretion and redistribution of metals to less sensitive target sites [29].

The results of the present and previous studies indicate that the toxicity of cadmium in amphibians is species dependent. The toxicity of cadmium correlates negatively with water hardness in aquatic organisms [43]. The toxicities of a particular toxicant to different species are influenced by factors such as temperature, pH, hardness and dissolve oxygen content of test water and physiological condition of the test animal [35].

CONCLUSION

The present study clearly indicates that cadmium caused mortality to the frog and this could be one of the reasons for population decline of frogs which inhabit water contaminated with heavy metals.

ACKNOWLEDGMENT

Authors are thankful to Dr. Sunil K. Srivastav for his help during the conduct of experiment. The authors declare that there is no conflict of interests.

REFERENCES

1. Ayenimo J, Yusuf A, Adekunle A, Makinde O. Heavy metal exposure from personal care products. Bull Environ Contam Toxicol 2010;84(1):8-14.
2. Gebhardt R. Prevention of cadmium-induced toxicity in liver-derived cells by the combination preparation Hepeel. Environ Toxicol Pharmacol 2009;27(3):402-9.
3. Chora S, Starita-Geribaldi M, Guignonis J-M, Samson M, Roméo M, Bebianno MJ. Effect of cadmium in the clam *Ruditapes decussatus* assessed by proteomic analysis. Aquat Toxicol 2009;94(4):300-8.
4. Järup L, Åkesson A. Current status of cadmium as an environmental health problem. Toxicol Appl Pharmacol 2009;238(3):201-8.
5. Wilson L. Cadmium: The pseudo-masculine mineral. 2008.
6. ATSDR. CERCLA priority list of hazardous substances. Atlanta, GA: Agency for Toxic Substances and Disease Registry, US Department of Health and Human Services. 2007.
7. Siu ER, Mruk DD, Porto CS, Cheng CY. Cadmium-induced testicular injury. Toxicol Appl Pharmacol 2009;238(3):240-9.
8. Goyer RA. Toxic and essential metal interactions. Annu Rev Nutr 1997;17(1):37-50.
9. Ye J-L, Mao W-P, Wu A-L, Zhang N-N, Zhang C, Yu Y-J, et al. Cadmium-induced apoptosis in human normal liver L-02 cells by acting on mitochondria and regulating Ca²⁺ signals. Environ Toxicol Pharmacol 2007;24(1):45-54.
10. Houlahan JE, Findlay CS, Schmidt BR, Meyer AH, Kuzmin SL. Quantitative evidence for global amphibian population declines. Nature 2000;404(6779):752-5.
11. Selvi M, Gül A, Yılmaz M. Investigation of acute toxicity of cadmium chloride (CdCl₂ · H₂O) metal salt and behavioral changes it causes on water frog (*Rana ridibunda* Pallas, 1771). Chemosphere 2003;52(1):259-63.
12. International Union for Conservation of Nature. Global Amphibian Assessment. International Union for Conservation of Nature, Conservation International, Nature Serve, Washington DC. USA. 2004.
13. Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues AS, Fischman DL, et al. Status and trends of amphibian declines and extinctions worldwide. Science 2004;306(5702):1783-6.
14. Blaustein AR, Hoffman PD, Hokit DG, Kiesecker JM, Walls SC, Hays JB. UV repair and resistance to solar UV-B in amphibian eggs: a link to population declines? P Natl A Sci 1994;91(5):1791-5.
15. Blaustein AR, Romansic JM, Kiesecker JM, Hatch AC. Ultraviolet radiation, toxic chemicals and amphibian population declines. Divers Distrib 2003;9(2):123-40.
16. Daszak P, Berger L, Cunningham AA, Hyatt AD, Green DE, Speare R. Emerging infectious diseases

- and amphibian population declines. *Emerg Infect Dis* 1999;5(6):735-48.
17. Chambouvet A, Gower DJ, Jirků M, Yabsley MJ, Davis AK, Leonard G, et al. Cryptic infection of a broad taxonomic and geographic diversity of tadpoles by Perkinsea protists. *P Natl A Sci* 2015;112(34):E4743-E51.
 18. Sparling DW, Lowe TP. Metal concentrations of tadpoles in experimental ponds. *ENVIRON POLLUT* 1996;91(2):149-59.
 19. Relyea RA. The lethal impact of Roundup on aquatic and terrestrial amphibians. *ECOL APPL* 2005;15(4):1118-24.
 20. Hayes TB, Case P, Chui S, Chung D, Haeffle C, Haston K, et al. Pesticide mixtures, endocrine disruption, and amphibian declines: are we underestimating the impact? *Environ Health Persp* 2006;114:40.
 21. Kats LB, Ferrer RP. Alien predators and amphibian declines: review of two decades of science and the transition to conservation. *Divers Distrib* 2003;9(2):99-110.
 22. Boone MD, Semlitsch RD, Little EE, Doyle MC. Multiple stressors in amphibian communities: effects of chemical contamination, bullfrogs, and fish. *Ecol Appl* 2007;17(1):291-301.
 23. Laurance WF. Global warming and amphibian extinctions in eastern Australia. *Austral col* 2008;33(1):1-9.
 24. Sparling DW, Fellers GM, McConnell LL. Pesticides and amphibian population declines in California, USA. *Environ Toxicol Chem* 2001;20(7):1591-5.
 25. Bridges C, Dwyer F, Hardesty D, Whites D. Comparative contaminant toxicity: are amphibian larvae more sensitive than fish? *B Environ Contam Tox* 2002;69(4):562-9.
 26. McDiarmid R W, Mitchell J C. Diversity and distribution of amphibians and reptiles. In: Sparling DW, Linder G, Bishop CA. *Ecotoxicology of Amphibians and Reptiles*. Society of Environmental Toxicology and Chemistry. 2000.p.15-69.
 27. Wef AA. Standard methods for the examination of water and wastewater. Washington, DC. 1998.
 28. Ezemonye L, Enuneku A. Evaluation of acute toxicity of cadmium and lead to amphibian tadpoles (Toad: *Bufo maculatus* and frog: *Ptychadena bibroni*). *J Aquat Sci* 2005;20(1):33-8.
 29. Enuneku AA, Ezemonye LI. Acute toxicity of cadmium and lead to adult toad *Bufo maculatus*. *Asian J Biol Life Sci* 2012;1: 238-41.
 30. Woodall C, Maclean N, Crossley F. Responses of trout fry (*Salmo gairdneri*) and *Xenopus laevis* tadpoles to cadmium and zinc. *Comp Biochem Physiol C: Comp Pharmacol*. 1988;89(1):93-9.
 31. Slooff W, Baerselman R. Comparison of the usefulness of the mexican axolotl (*Ambystoma mexicanum*) and the clawed toad (*Xenopus laevis*) in toxicological bioassays. *Bull Environ Contam Toxicol* 1980;24(1):439-43.
 32. De Zwart D, Slooff W. Toxicity of mixtures of heavy metals and petrochemicals to *Xenopus laevis*. *Bull Environ Contam Toxicol* 1987;38(2):345-51.
 33. Ferrari L, Salibián A, Muiño CV. Selective protection of temperature against cadmium acute toxicity to *Bufo arenarum* tadpoles. *Bull Environ Contam Toxicol* 1993;50(2):212-8.
 34. Grillitsch B, Chovanec A. Heavy metals and pesticides in anuran spawn and tadpoles, water, and sediment. *Toxicol Environ Chem* 1995;50(1-4):131-55.
 35. Loubourdis N, Kyriakopoulou-Sklavounou P, Zachariadis G. Effects of cadmium exposure on bioaccumulation and larval growth in the frog *Rana ridibunda*. *Environ Pollut* 1999;104(3):429-33.
 36. McCarty L, Henry J, Houston A. Toxicity of cadmium to goldfish, *Carassius auratus*, in hard and soft water. *J Fish Res Board Can* 1978;35(1):35-42.
 37. Pickering QH, Henderson C. The acute toxicity of some heavy metals to different species of warmwater fishes. *Air Water Pollut* 1966;10(6):453-63.
 38. Benson WH, Baer KN, Stackhouse RA, Watson CF. Influence of cadmium exposure on selected hematological parameters in freshwater teleost, *Notemigonus crysoleucas*. *Ecotoxicol Environ Saf* 1987;13(1):92-6.
 39. Akram M, Hafeez M, Nabi G. Histopathological changes in the kidney of a freshwater cyprinid fish, *Barilius vagra*, following exposure to cadmium. *Pak J Zool* 1999;31(1):77-80.
 40. Dutta TK, Kaviraj A. Acute toxicity of cadmium to fish *Labeo rohita* and copepod *Diaptomus forbesi* pre-exposed to CaO and KMnO₄. *Chemosphere* 2001;42(8):955-8.
 41. Rai R, Mishra D, Srivastav SK, Srivastav Ajai K. Acute toxicity of cadmium against catfish, *Heteropneustes fossilis* (Siluriformes: Heteropneustidae) in static renewal bioassays. *Ethiopian J Biol Sci* 2008; 7: 185-91.
 42. Medina MF, Cosci A, Cisint S, Crespo CA, Ramos I, Villagra ALI, et al. Histopathological and biological studies of the effect of cadmium on *Rhinella arenarum* gonads. *Tissue and Cell* 2012;44(6):418-26.
 43. USEPA. 2001 Update of Ambient Water Quality Criteria for Cadmium. Environmental Protection Agency Washington, DC; 2001.