

**Original Article****Histological Effects of Cadmium on Hepatopancreas and Gill in *Cyprinus carpio***Farzad Ghiasi\*<sup>1</sup>, Seyed Saeed Mirzargar<sup>2</sup>, Javad Ashrafihellan<sup>3</sup>

Received: 01.03.2017

Accepted: 05.04.2017

**ABSTRACT**

**Background:** Histological studies in organs like hepatopancreas and gill of the freshwater fish, *Cyprinus carpio*, were made to assess tissue damage due to sub-lethal concentration of cadmium after a long exposure period.

**Methods:** This study was conducted in University of Tehran, in 2013. The fish were divided into two groups in 1000 litre fiberglass tanks supplied with dechlorinated water [hardness 302.6 mg CaCO<sub>3</sub>/l, pH 7, O<sub>2</sub> 7.8 mg/L and temperature 15 ± 2 °C] and continuous aeration. Group 1 was without any cadmium considered as control group. Group 2 was exposed to 30 ppb CdCl<sub>2</sub> (Merck) containing 7.8 ppb cadmium. The tissue samples from: hepatopancreas, gill were collected 30 days post exposure and processed by histological procedures.

**Results:** The main lesions in cadmium exposed groups were: (a) pancreatitis necrosis in endocrine part of pancreas (b) cholangitis and necrosis in liver parenchyma (c) hypertrophy, fusion and telangiectasia in secondary lamellae

**Conclusion:** Cadmium chloride at low concentration can induce pathological alterations in hepatopancreas and gill of common carp.

**Keywords:** Cadmium, Common Carp, Gill, Hepatopancreas, Histological Effects.

IJT 2017 (5): 1-4

**INTRODUCTION**

Cadmium is a non-essential element that can have severe toxic effect on aquatic organisms when present in excessive amount. Cadmium is important because of its use in various industrial processes, as by-product of zinc mining, fossil fuel, base metal smelting and combustion [1]. The role of sub-lethal concentration of cadmium in changing haematological parameters in *Hetroclarias* is reported [2] and lysozyme, leukocyte count and phagocytic index in *Cyprinus carpio* [3]. There are many investigations on the histopathological changes in common carp kidney due to cadmium exposure [4, 5]. Randi et al. [1] studied the histopathological effects of cadmium on the gill of *Macropsobrycon uruguayanae*. Pathological damages in the liver of *Oreochromis mossambicus* after short and long time exposure to cadmium and zinc was documented [6]. Histopathological alterations of white sea bass, *Lates calcarifer* in acute and sub-chronic cadmium exposure have been reported [7].

Exposures to low levels of cadmium can cause DNA damage and stress in common carp [8]. Cadmium is a risk factor for pancreatic cancer in human [9].

The goal of the current study was to identify whether cadmium at sub-lethal concentration, can cause histological changes in the hepatopancreas and gill of *C. carpio*.

**MATERIALS AND METHODS**

The current study was done in University of Tehran in 2013. Common carp obtained from a local fish farm were acclimated in holding tank for 1 week, then 20 apparently healthy fish, mean weight 700 g were randomly divided into two groups (1-2) in 1000 litre indoor fiberglass tanks supplied with dechlorinated water (hardness 302.6 mg/L as CaCO<sub>3</sub>, dissolved oxygen 7.8 mg/L, pH 7, temperature 15 ± 2°C), without water flow and with continuous aeration. Fish in group 1 was held in water without cadmium and group 2 exposed to 30 ppb cadmium chloride (Merck) for 30 d This

1. Department of Fisheries, Faculty of Natural Resources, University of Kurdistan, Sanandaj, Iran.

2. Department of Aquatic Animal Health, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran.

3. Department of Pathology, Veterinary college, University of Tabriz, Tabriz, Iran.


\*Corresponding Author E-Mail: f.ghiasi@uok.ac.ir

concentrations were less than 1% of LC50 96 h for this species [10] [11]. Water quality was measured throughout the exposure two times weekly. The fish were fed 1% body weight weekly with commercial food. Five fish from each tank were euthanized and sampled for tissues material containing hepatopancreas, gill 30 d post exposure. Tissue samples were fixed in 10% buffered formalin, embedded in paraffin, Sectioned at  $5\mu$ , and stained with hematoxylin and eosin (H&E) for histopathology [12].


All procedures were conducted according to ethical guidelines of animal experiences (CEE 86/609 regulation).

## RESULTS


No death occurred either in the control or in the experimental group during the whole period of the experiment. After 30 days post challenge histopathological alterations in group 2 were as follows: Focal necrosis and fatty change (Fig. 1) and mononuclear cholangitis (Fig. 2,3) were seen in the liver.



**Figure 1.** Focal necrosis (arrow) and fatty change in the liver (H&E  $\times 400$ ).




**Figure 2.** Mononuclear cholangitis in the liver (arrow (H&E  $\times 200$ )).




**Figure 3.** Mononuclear cholangitis in the liver (arrow) (H&E  $\times 400$ ).


Mononuclear (lymphocytic) pancreatitis (Fig. 4) and single cell necrosis were observed in pancreas (Fig. 5). Hyperplasia of secondary lamellae, lamellar hypertrophy and fusion and lamellar clubbing (Fig. 6) and telangiectasia in the secondary gill lamellae (Fig. 7) were seen. No histopathological changes were observed in control group during experiments period.




**Figure 4.** Necrosis in the endocrine portion of pancreas (arrow)(H&E  $\times 200$ ).



**Figure 5.** Pancreatitis and necrosis in the endocrine portion of pancreas (arrow) (H&E  $\times 200$ ).



**Figure 6.** Hyperplasia of secondary lamellae, lamellar hypertrophy and fusion and lamellar clubbing (H&E×200).



**Figure 7.** Telangiectasia in the secondary lamellae (arrow) (H&E×200).

## DISCUSSION

Histology supplies a fast way to detect effects of irritants, especially chronic ones, in various tissues and organs [13]. In the current study, tissue abnormalities were found in all investigated organs. As our findings after 30 d post challenge telangiectasia in gills, focal necrosis in endocrine part of pancreas, fatty changes and focal necrosis in liver were only observed in cadmium exposure group. The cadmium induced histopathological changes in the gill of *C. carpio* were similar to lesions shown in other fish due to heavy metal toxicity [1]. Our findings are supported by previous finding that showed fusion of adjacent secondary lamellae and lamellar telangiectasia were only induced by exposure to cadmium while hyperplasia of secondary lamellar epithelium produced by either cadmium, or starvation even exposure time [1].

Cadmium chloride at sub-lethal dose can cause hypertrophy of lamellar epithelium, destroy of gill lamellar, blood congestion, hypertrophy and necrosis in liver of fresh water fish

*Ophicephalus striatus* [14]. Necrosis in liver induced by cadmium in fresh water fish *Channa striatus* is reported [15]. In tissues such as kidney, spleen, liver and gill a low concentration of cadmium can be enlarged [16]. Seven percent of the human populations have kidney dysfunction due to cadmium exposure [17]. Metallothioneins (MTs) are cysteine rich proteins with low-molecular-weight, which can bind with metal ions [18]. It is critical for protecting human health from cadmium toxicity [17]. It is an important protein in the cellular defense against Cd toxicity and lethality [19]. MT synthesis in fish is related with organs involved in metal uptake, metabolism, and excretion, such as gill, liver, kidney, and intestine [20].

Possible ways by which MT may protect against metal toxicity include reduction of metal transport into cells; sequestration of metal within the cells; and increasing metal export out of cells. The mechanism of MT in cellular defense against Cd toxicity includes removal of free (toxic) cadmium from the inactive Cd-MT complex and alter the intracellular distribution of Cd [19] pathological effects would occur when the rate of metal uptake exceeds the rate of metallothionein. Once the metallothionein pool is saturated, free cadmium initiate toxic damage to the cells.

## CONCLUSION

Cadmium chloride at low concentration can cause pathological damage in hepatopancreas and gill of common carp. Tissue abnormalities were found in all investigated organs. Pathological injury and telangiectasia of gill filaments were found. Liver tissue revealed hemorrhages, blood congestion and necrotic cells with mononuclear cell infiltration, but for clear detection of the damages we need to use quantitative histopathology methods. Histopathology could be used as an effective index to monitor toxicological changes induced by polluted environment and fish health status during exposure to pollutants such as heavy metals.

## ACKNOWLEDGEMENT

We would like to thank Miss Andisheh Kaseb for helping us. The authors declare that there is no conflict of interest.

## REFERENCES

1. Randi A, Monserrat J, Rodriguez E, Romano L. Histopathological effects of cadmium on the gills

- of the freshwater fish, *Macropsobrycon uruguayanae* Eigenmann (Pisces, Atherinidae). J Fish Dis 1996;19(4):311-22.
2. Kori-Siakpere O, Ake J, Avworo U. Sublethal effects of cadmium on some selected haematological parameters of *Heteroclinus* (a hybrid of *Heterobranchius bidorsalis* and *Clarias gariepinus*). Int.j.zool.res 2006;2(1):77-83.
  3. Ghiasi F, Mirzargar S, Badakhshan H, Shamsi S. Effects of low concentration of cadmium on the level of lysozyme in serum, leukocyte count and phagocytic index in *Cyprinus carpio* under the wintering conditions. J Fisheries Aqua Sci 2010;5(2):113-9.
  4. Ghiasi F, Mirzargar S. Pathological effects of cadmium and efficiency of natural zeolite, Clinoptilolite, to reduce the cadmium toxicity in common carp and its ability to remove Cadmium from contaminated water. Res J Vet Pract 2015;3(2):36-40.
  5. Singhal R, Jain M. Cadmium-induced changes in the histology of kidneys in common carp, *Cyprinus carpio* (Cyprinidae). B Environ Contam Toxicol 1997;58(3):456-62.
  6. Van Dyk JC, Pieterse G, Van Vuren J. Histological changes in the liver of *Oreochromis mossambicus* (Cichlidae) after exposure to cadmium and zinc. Ecotoxicol Environ Saf 2007;66(3):432-40.
  7. Thophon S, Kruatrachue M, Upatham E, Pokethitiyook P, Sahaphong S, Jaritkhuan S. Histopathological alterations of white seabass, *Lates calcarifer*, in acute and subchronic cadmium exposure. Environ Pollut 2003;121(3):307-20.
  8. Jia X, Zhang H, Liu X. Low levels of cadmium exposure induce DNA damage and oxidative stress in the liver of Oujiang colored common carp *Cyprinus carpio* var. color. Fish Physiol Biochem 2011;37(1):97-103.
  9. Schwartz GG, Reis IM. Is cadmium a cause of human pancreatic cancer? Cancer Epidemiol Biomarkers Prev 2000;9(2):139-45.
  10. Suresh A, Sivaramakrishna B, Radhakrishnaiah K. Patterns of cadmium accumulation in the organs of fry and fingerlings of freshwater fish *Cyprinus carpio* following cadmium exposure. Chemosphere 1993;26(5):945-53.
  11. Witeska M, Kondera E, Szczygielska K. The effects of cadmium on common carp erythrocyte morphology. Pol J Environ Stud. 2011;20:783-8.
  12. Bernet D, Schmidt H, Meier W, Burkhardt-Holm P, Wahli T. Histopathology in fish: proposal for a protocol to assess aquatic pollution. J Fish Dis 1999;22(1):25-34.
  13. Johnson LL, Stehr CM, Olson OP, Myers MS, Pierce SM, Wigren CA, et al. Chemical contaminants and hepatic lesions in winter flounder (*Pleuronectes americanus*) from the Northeast Coast of the United States. Environ Sci Technol 1993;27(13):2759-71.
  14. Bais U, Lokhande M. Effect of cadmium chloride on histopathological changes in the freshwater fish *Ophiocephalus striatus* (Channa). Int J Zoological Res 2012;8(1):23-32.
  15. Radhakrishnan M, Hemalatha S. Sublethal toxic effects of cadmium chloride to liver of freshwater fish *Channa striatus* (Bloch). Ame-Euras J Toxicol Sci 2010;2(1):54-6.
  16. Ghiasi F, Mirzargar S, Badakhshan H, Amoli J. Influence of Iranian natural zeolite on accumulation of cadmium in *cyprinus carpio* tissues following exposure to low concentration of cadmium. Asian J Animal Vet Adv 2011;6(6):636-41.
  17. Klaassen CD, Liu J, Diwan BA. Metallothionein protection of cadmium toxicity. Toxicol Appl Pharmacol 2009;238(3):215-20.
  18. Nordberg GF. Cadmium and health in the 21st century—historical remarks and trends for the future. Biometals 2004;17(5):485-9.
  19. Park JD, Liu Y, Klaassen CD. Protective effect of metallothionein against the toxicity of cadmium and other metals. Toxicol 2001;163(2):93-100.
  20. Mieiro C, Bervoets L, Joosen S, Blust R, Duarte A, Pereira M, et al. Metallothioneins failed to reflect mercury external levels of exposure and bioaccumulation in marine fish—Considerations on tissue and species specific responses. Chemosphere 2011;85(1):114-21.