

Accepted Manuscript**Accepted Manuscript (Uncorrected Proof)**

Title: Evaluation of Clinical and Paraclinical Manifestations of Mushroom Poisoning: A Cross-Sectional Study

Running Title: Manifestations of Mushroom Poisoning

Authors: Ghafarali Mahmoudi¹, Mohammad Hojat Bazrafkan², Golnaz Mahmoudvand², Peyman Astaraki¹, Soudabeh Zare³, Arian Karimi Rouzbahani^{2,*}

1. *Department of Internal Medicine, Faculty of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran.*
2. *Student Research Committee, Lorestan University of Medical Sciences, Khorramabad, Iran.*
3. *Department of Biostatistics and Epidemiology, School of Health and Nutrition, Lorestan University of Medical Sciences, Khorramabad, Iran.*

***Corresponding Author:** Arian Karimi Rouzbahani, Student Research Committee, Lorestan University of Medical Sciences, Khorramabad, Iran. Email: ariankarimi1998@gmail.com

To appear in: **Iranian Journal of Toxicology**

Received date: 2022/08/05

Revised date: 2022/09/27

Accepted date: 2022/10/03

This is a “Just Accepted” manuscript, which has been examined by the peer-review process and has been accepted for publication. A “Just Accepted” manuscript is published online shortly after its acceptance, which is prior to technical editing and formatting and author proofing. *Iranian Journal of Toxicology* provides “Just Accepted” as an optional and free service which allows authors to make their results available to the research community as soon as possible after acceptance. After a manuscript has been technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as a published article. Please note that technical editing may introduce minor changes to the manuscript text and/or graphics which may affect the content, and all legal disclaimers that apply to the journal pertain.

Please cite this article as:

Mahmoudi G, Bazrafkan MH, Mahmoudvand G, Astaraki P, Zare S, Karimi Rouzbahani A. (In Press). Evaluation of Clinical and Paraclinical Manifestations of Mushroom Poisoning: A Cross-Sectional Study. *Iranian Journal of Toxicology*. Just Accepted publication Aug. 16, 2022. Doi: <http://dx.doi.org/10.32598/IJT.17.1.1018.1>

DOI: <http://dx.doi.org/10.32598/IJT.17.1.1018.1>

ABSTRACT

Background: Mushroom poisoning is a major health condition with a wide range of clinical and paraclinical features. This study aimed at evaluating the frequency of clinical and paraclinical manifestations of mushroom poisoning in patients referred to Shahid Rahimi Hospital in Khorramabad, Iran, over a one-year period (2018-2019).

Methods: The data collected were associated with the clinical manifestations, age, sex, seasons, type of mushrooms, patients' residence, latent phase, clinical and laboratory findings, length of hospital stay, interventions and the treatments. The underlying diseases were also recorded. After data collection, they were entered into SPSS, version 18 software and analyzed statistically.

Results: 124 patients with a mean age of 36.65 years old were recruited into the study, 73 of whom were male and 51 female. The mean duration of the hospital stay was 2.19 days. The mean time elapsed between the consumption and the symptoms development was 4.42 hours. Similarly, the duration between the consumption and referral to the hospital was 4.72 hours. Most cases occurred in the Spring (91.1%). The most common clinical signs in the poisoned subjects were nausea and vomiting (81.5%). The most therapeutic medications were Livergol (48.4%) and Atropin (33.1%), and most subjects had consumed mushrooms grown in the nature (79.8%). One person died because of the poisoning (0.8%).

Conclusions: A large majority of the patients developed nausea and vomiting, whom were treated with drugs, but one patient died. People should be aware of, warned against, and educated about the types of mushrooms before consumption.

Keywords: Amanita mushrooms; Clinical symptoms; Mushroom poisoning; Nausea; Vomiting

INTRODUCTION

Some mushrooms may be nutritious foods [1], and it has been estimated that 1.5 million different species of them exist worldwide [2]. However, due to the existence of poisonous mushrooms, it's crucial to identify the type before consumption [1]. Poisonous mushrooms contain toxic substances, such as amatoxins, muscarine, coprine, orellanus, and psilocybin, that are hazardous to human health [3, 4]. They are subdivided into groups based on the clinical signs and symptoms they cause in humans at presentation. Common syndromes associated with mushroom poisoning include damage to major internal organs, neurotoxicity, metabolic, endocrine and gastrointestinal disorders [5]. Most poisoning cases present with gastroenteritis and psychoneurological disorders; however, the patients usually recover without serious clinical consequences [6]. The most dangerous fungal species known today are Amanita mushrooms (*A. phalloides*, *A. virosa* & *A. verna*), which cause severe clinical symptoms or even fatal outcome [7].

Three stages have been described for Amanita poisoning: *a*) gastrointestinal symptoms in the first 6 to 24 hours, *b*) hepatic sign and symptoms appear with a 12-48 hour latency, and *c*) acute kidney and liver failure develop 24-72 hours later [8]. Although certain methods have been suggested to identify toxic mushrooms, there is no definitive standards for their identification [4]. Features such as the mushroom's shape, surface, and color of the cap, gill, stalk, and the odor might help differentiate safe mushrooms from the poisonous ones [9]. The prevalence of mushroom toxicity has been estimated at 7428 cases per year in the United States [10]. In Iran, a total of 50 species of poisonous mushrooms have been identified. In the Spring, sporadic cases of mushroom poisoning occur in Iran [11]. In a study at a poisoning center in Tehran, 37 cases of mushroom poisoning were reported over a 10-year period [12].

In the diagnostic process, the clinical differences for various poisoning syndromes should be distinguished [5]. Identification of the consumed mushrooms and the time elapsed between the consumption and onset of symptoms are the key to the diagnosis. Macroscopic identification of the ingested mushrooms or the leftovers, and the microscopic features of the spores in the plant are the most useful data for the diagnosis [8]. To date, such a study has not been conducted.

Aim of the Study: Khorramabad city is located in Lorestan province, a mountainous area in western Iran. Due to its unique geographical nature, a high variety of herbal plants is grown in this area, thus leading to a high incidence poisoning [13]. In 2018, the second highest mushroom poisoning outbreak in Iran occurred in Lorestan province [11]. Given the high rate of mushroom poisoning reported in Lorestan province, this study aimed to evaluate the frequency of clinical and paraclinical manifestations of poisoning cases due to the consumption of mushrooms, as referred to Shahid Rahimi Hospital in Khorramabad, Iran, during a one year period (2018-2019).

MATERIALS and METHODS

Study Design and Participants: This cross-sectional, descriptive study was undertaken to investigate the frequency of clinical and paraclinical manifestations due to mushroom poisoning at Shahid Rahimi Hospital in Khorramabad, Iran. The study subjects were patients who were referred to, diagnosed and hospitalized due to mushroom poisoning. Patients with incomplete medical records and those who did not consent to participate in the study were excluded.

Data Collection: The frequency of clinical manifestations of mushroom poisoning, such as gastrointestinal or neurological symptoms, and others, such as the age, sex, seasons, type of mushrooms consumed, patient's residence, latent phase, clinical and laboratory findings, length of hospital stay, therapeutic interventions, and the underlying diseases were reviewed and recorded. Initially, the responsible factors were tabulated as a checklist of full description and review of the each useful study found over a literature search.

Data Analyses: After collecting the relevant data and clinical information, descriptive statistical methods were used to analyze the data. A statistical software (SPSS, version 18) was utilized to analyze the data. Using descriptive methods, the frequency, percentage, mean and standard deviation, and other findings are presented in this article as one Figure and nine Tables.

Ethical Considerations: The study protocol was approved by the Research Ethics Committee at Lorestan University of Medical Sciences (Code #: IR.LUMS.REC.1399.064). The data for each patient were collected from the available clinical records, such that there was no direct communication with the patients. They had signed a consent form at admission into the hospital. All patients' information were kept strictly confidential and no disclosure was made to any sources other than the attending physicians. In order to avoid registering the patients by their names, we used a specific code to refer to each individual. The information was kept confidential and recorded in numerical codes, using SPSS software, version 18. Prior to conducting this study, necessary arrangements were made with the administration of the Lorestan University of Medical Sciences' Hospital via submitting a formal document, containing the study protocol and the aims. Subsequently, the Ethics Committee reviewed the study protocol and approved it.

RESULTS

In this study, 124 patients with mushroom poisoning participated, 73 of whom (58.9%) were male and 51 (41.1%) were female. The patients' mean age was 36.65 (± 19.7) years old, with the minimum being 2 and the maximum 83 years old. Figure 1 represents the study population age distribution. The mean duration of hospital stay was 2.19 (± 1.7) days, ranging from zero to seven days. In terms of patients' occupation, the highest frequency of the study population were housewives (36.3%). Most of the poisoning cases occurred in the Spring (91.1%), and the most frequently used medications to treat the symptoms and side effects were Livergol (48.4%), and Atropin (33.1%). See Table 1.

The mean and standard deviation for the duration of the mushroom consumption until being symptomatic was 4.58 hours, ranging from zero to 24 hours. The duration from the mushroom consumption to the patients being referred to the hospital was 6.05 hours (minimum, 0.5, maximum 24 hours). Table 2 lists the data for the mean, standard deviation, minimum and maximum laboratory findings.

Most of the subjects had collected the mushrooms from the nature (79.8%). As seen in Table 3, the most common clinical symptoms reported by the patients were nausea and vomiting (81.5%), sweating (41.1%), and abdominal pain (37.1%). Of the total cases, 23 of them (18.5%) were examined by a second physician before being admitted to the hospital. The most frequent underlying diseases were as follows: hypertension (8.9%); coronary artery disease (6.5%); diabetes (5.6%); renal disease (1.6%); hematological disease (1.6%); and liver disease (0.8%). Considering the final outcomes, 75 patients (60.5%) were hospitalized, while 36 of them (29%)

left the hospital voluntarily, 9 (7.3%) were admitted to the ICU, 2 (1.6%) were discharged from emergency room, 1 (0.8%) was referred to a specialized center, and one patient (0.8%) died. Table 4 presents the distribution of the laboratory findings for the subjects.

Table 5 provides information on the distribution of the subjects' clinical signs based on gender.

Table 6 lists the data on the distribution of clinical signs based on the patients' occupation.

Table 7 presents information on the clinical signs for each patient based on the season.

Table 8 shows the information on the clinical signs based on the patients' underlying diseases.

Finally, Table 9 presents the patients' signs based on the clinical outcomes.

DISCUSSION

This study evaluated the clinical and paraclinical manifestations observed in patients with mushroom poisoning in Khorramabad, Lorestan province, Iran. In Iran, a total number of 50 species of poisonous mushrooms have been identified and *Amanita* is the most dangerous species identified in the country [14]. In the 2018 mushroom poisoning outbreak in Iran, the following species were involved: *Lepiota brunneioncarnata*, *Hypholoma fascicalare*, and *Coprinopsis atramentaria* [11]. In a previous study in Lorestan province, muscarin toxin was reported as the most common cause of mushroom poisoning [13].

Demographic Characteristics: Regarding the demographic characteristics of patients, most of our subjects were male. Consistent with our findings, Chen, *et al.* [15] and Pajoumand, *et al.* [12] also observed a higher incidence of poisoning in men. The mean age of our patients was 36.65, consistent with 35 year old as reported by Erenler, *et al.* [17]. The mean length of hospital stay in our study was 2.19 days. However, in another study by Keller, *et al.* [18], 78.4% of cases stayed in the ER for less than 24 hours, similar to 1.89 days that was observed by Khatir, *et al.* [19].

Our findings showed that 79.8% of the cases had consumed self-picked poisonous mushrooms, which is similar to 83% of the patients, that was found by Jiang, *et al.* [20]. Further, we found the mean time elapsed between the mushroom consumption and the onset of symptoms was 4.58 hours. Also, the interval between the mushroom consumption and the admission to the hospital was 6.05 hours. Our findings were similar to those reported by two earlier studies [18, 20] who observed that in 68.6% and 62% of the subjects, respectively, the clinical symptoms began within the first 6 hours of the mushroom ingestion. Also, with respect to the season of the year when the poisoning happened, we found that 91.1% of them occurred in the Spring. Similarly, Varshouchi, *et al.* reported that 68% of poisoning cases occurred in the Spring [21]. However, Gold, *et al.* [22] have reported that a greater number of similar poisoning (36.8%) happened in the Summer while only 22.5% of them occurred in the Spring.

Clinical and Paraclinical Findings: In our study, the most frequently reported clinical symptoms by the patients were nausea and vomiting, followed by sweating, abdominal pain, and diarrhea. Nausea and vomiting was the most common symptom in both male and female, and in patients with various underlying diseases. However, in retired patients and employees, and in those who were admitted in the Fall, sweating and abdominal pain were the most frequently reported symptoms. Similarly in 2016, Chan, *et al.* reported 67 mushroom poisoning cases in Hong Kong, 90% of whom presented with gastrointestinal symptoms [16]. In these patients, all of the laboratory tests were within normal ranges. Evidently, the identification of the type of mushrooms consumed and the knowledge of clinical variations among poisoning syndromes

remain the major diagnostic challenge to date [5, 8]. However, a significantly higher neutrophil-to-lymphocyte ratio and lower platelet count have been observed in patients with mushroom poisoning [23]. In our study population, impaired liver function test was the most common paraclinical finding. Similar observations have also been reported in two previous studies performed in Iran [14, 24].

Therapeutic Interventions: Currently, there is no definitive and preferred treatment for the human cases with mushroom poisoning. Although there are controversies over the use of activated charcoal, this approach has been recommended by many earlier studies. Supportive treatments, such as correction of electrolyte abnormalities and dehydration should also be considered [25]. Ordinarily, drugs and antidotes are used in some poisoning cases. In case of a neurological sign, such as seizure, benzodiazepines and pyridoxine may be prescribed. Also, atropine is beneficial to counter the cholinergic presentations. In cases of Amanita toxicity, patients may benefit from Silibinin, N-acetylcysteine, or even liver transplant if a fulminant hepatic failure has occurred [18].

In our study, the most common therapeutic interventions were Livergol (containing *Silybum marianum*), atropine, and N-acetylcysteine. In Keller, *et al.*'s study, 56.9% of the cases received activated charcoal, and the other 24% were treated with N-acetylcysteine [18]. The overall mortality rate of mushroom poisoning cases is fairly low at 2.17% [6], although, a wide range of mortality has been reported in the reviewed literature. In our study, this rate was very low, at 0.8% of the mushroom poisoning cases. Chen, *et al.* reported only two deaths among the 429 patients they treated with mushroom poisoning in China [15]. Conversely, De Olano, *et al.* have reported a higher rate of mortality at 8.8% among their poisoning cases that ingested cyclopeptide mushrooms in the United States [26].

Limitations of the Study: We had the following limitations in this study: *a*) the medical records of some patients were incomplete, which led to their exclusion from the study; *b*) the types of consumed mushrooms had not been recorded in all patients record.

CONCLUSIONS

In this study, a large majority of the patients developed gastrointestinal (GI) symptoms, such as nausea and vomiting, whom were managed clinically with medications; however, there was one fatal case. In addition to the GI symptoms, our subjects complained of sweating and abdominal pain. Among them, the most frequent underlying diseases were hypertension, coronary artery disease, diabetes, and renal, hematological or liver diseases. Health authorities are highly advised to educate the public on poisoning due to mushroom ingestion. Such efforts will positively lead to reductions the number of poisoning cases, especially in the Spring and Summer. It will also shorten the patients' hospital stay, and ultimately reduce the subsequent mortality.

Conflict of Interest: The authors declare no conflict of interest with any entities to disclose.

Compliance with Ethical Guideline: This study was approved by the Research Ethics Committee at Lorestan University of Medical Sciences (Ethics code: IR.LUMS.REC.1399.064).

Funding: This study did not receive funds from any entity other than the authors.

Authors' Contributions: Dr. GA Mahmoudi: conceptualized and designed the study, and reviewed and revised the manuscript. Dr. MH Bazrafkan: collected the data. Dr. G Mahmoudvand: drafted the initial manuscript. Dr. P Astaraki: provided professional consultation in support of the study conduction. Dr.S Zare: designed the data collection instruments and carried out the data analyses. Mr. A. Karimi Rouzbahani: coordinated and supervised data collection, and critically reviewed the manuscript for its intellectual contents. All authors read and approved the final version of the manuscript prior to submission to the Journal.

Acknowledgment: The authors gratefully appreciate the participation of the patients in this cross-sectional clinical trial. The authors are also thankful to the administration and staff of Shahid Rahimi Hospital in Khorramabad, Iran, for their generous cooperation with this research project.

Accepted Manuscript (Uncorrected Proof)

REFERENCES

1. Zahan, N., *et al.*, A Deep Learning-Based Approach for Edible, Inedible and Poisonous Mushroom Classification. *2021 International Conference on Information and Communication Technology for Sustainable Development (ICICT4SD)*. 2021; 440-444.
2. Chitayae, N. and A. Sunyoto, Performance Comparison of Mushroom Types Classification Using K-Nearest Neighbor Method and Decision Tree Method. *3rd International Conference on Information and Communications Technology (ICOIACT)*. 2020; 308-313.
3. Sevindik, M., *Poisonous mushroom (nonedible) as an antioxidant source*. *Plant Antioxidants and Health*. 2020; 1-25.
4. Jo, W.-S., M.A. Hossain, and S.-C. Park, Toxicological profiles of poisonous, edible, and medicinal mushrooms. *Mycobiology*. 2014; 42(3): 215-220.
5. White, J., *et al.*, Mushroom poisoning: A proposed new clinical classification. *Toxicon*. 2019; 157: 53-65.
6. Li, H., *et al.*, Mushroom Poisoning Outbreaks— China, 2021. *China CDC Weekly*. 2022; 4(3): 35-40.
7. Tavassoli, M., *et al.*, Toxicological profile of Amanita virosa – A narrative review. *Toxicology Reports*. 2019; 6:143-150.
8. Wennig, R., *et al.*, Mushroom Poisoning. *Deutsches Arzteblatt International*. 2020;117(42):701-708.
9. Ismail, S., A.R. Zainal, and A. Mustapha, Behavioural features for mushroom classification. 2018 *IEEE Symposium on Computer Applications & Industrial Electronics (ISCAIE)*. 2018;412-415.
10. Brandenburg, W.E. and K.J. Ward, Mushroom poisoning epidemiology in the United States. *Mycologia*. 2018; 110(4):637-641.
11. Soltaninejad, K., Outbreak of Mushroom Poisoning in Iran: April-May, 2018. *The International Journal of Occupational and Environmental Medicine*. 2018; 9(3):152-156.
12. Pajoumand, A., *et al.*, A retrospective study of mushroom poisoning in Iran. *Human & Experimental Toxicology*. 2005; 24(12):609-613.
13. Mahmoudi, G.A., H. Solhi, and M.J. Tarrahi, The Study of Poisoning Whit Mushroom and Toxic Plants (Tattore and Ricinus Communis) Incidence in Patients Who Admitted in Shohadaie Ashaier and Madani Hospitals in the Spring and Summer of 2007. *Iranian Journal Toxicol*. 2007; 1(3): 4-4.
14. Dadpour, B., *et al.*, Mushroom poisoning in the northeast of Iran; a retrospective 6-year epidemiologic study. *Emergency (Tehran, Iran)*. 2017; 5(1): e23-e23.
15. Chen, L., *et al.*, Epidemiological analysis of wild mushroom poisoning in Zhejiang province, China, 2016–2018. *Food Science & Nutrition*. 2022;10(1): 60-66.
16. Chan, C., *et al.*, Mushroom poisoning in Hong Kong: a ten-year. *Hong Kong Med Journal*. 2016; 22(2): 124-130.
17. Erenler, A.K., *et al.*, Investigation of toxic effects of mushroom poisoning on the cardiovascular system. *Basic, Clinical Pharmacology & Toxicology*. 2016;119(3): 317-321.
18. Keller, S.A., *et al.*, Mushroom poisoning—A 17 year retrospective study at a level I university emergency department in Switzerland. *International Journal of Environmental Research and Public Health*. 2018; 15(12): 2855.
19. Khatir, I., *et al.*, Demographic and epidemiologic evaluation of mushroompoisoning: a retrospective study in 4-year admissions of Razi Hospital (Qaemshahr, Mazandaran, Iran). *Medicinski Glasnik: Official Publication of the Medical Association of Zenica-doboj Canton, Bosnia and Herzegovina*, 2020; 17(1): 117-122.
20. Jiang, L., M. Luo, and L. Hao, Epidemiological characteristics of mushroom poisoning in Yunnan Province, China, 2004-2016. *Southeast Asian Journal of Tropical Medicine and Public Health*. 2018; 49(3): 509-515.
21. Varshochi, M. and B. Naghili, Mushroom poisoning in northwest of Iran. *Iranian Journal of Clinical Infectious Diseases*. 2007; 2(4): 169-175.

22. Gold, J.A.W., *et al.*, Health Care Utilization and Outcomes Associated with Accidental Poisonous Mushroom Ingestions - United States, 2016-2018. *MMWR. Morbidity and Mortality Weekly Report*. 2021; 70(10): 337-341.
23. Doğan, M. and İ. Karahan, Evaluation of complete blood count parameters, cell ratios, and cell volume indices in mushroom poisonings. *Human & Experimental Toxicology*. 2019;38(10):1127-1131.
24. Badsar, A., *et al.*, Mushroom Poisoning in the Southwest Region of the Caspian Sea, Iran: A Retrospective Study. *Iranian Journal of Toxicology*. 2013; 7(20):798-803.
25. Beaumier, M., *et al.*, Mushroom poisoning presenting with acute kidney injury and elevated transaminases. *Kidney International Reports*. 2019; 4(6):877-881.
26. De Olano, J., *et al.*, Current fatality rate of suspected cyclopeptide mushroom poisoning in the United States. *Clinical Toxicology*. 2021; 59(1): 24-27.

Accepted Manuscript (Uncorrected Proof)


FIGURE

Figure 1: Histogram of the subjects' age distribution.

TABLES**Table 1:** Characteristics of the study subjects.

Category	Occupation	Frequency	Percent
Study population	Housewife	45	36.3
	Self-employed	34	27.4
	Unemployed	21	16.9
	Employees	7	5.6
	Students	6	4.8
	Retired	4	3.2
	Farmers	3	2.4
	Workers	2	1.6
	University students	2	1.6
	Total	124	100
Poisoning season	Spring	113	91.1
	Summer	6	4.8
	Fall	2	1.6
	Winter	3	2.4
Treatment	Livergol	60	48.4
	Atropine	41	33.1
	NAC	17	13.7
	Charcoal	4	3.2
	FFP	1	0.8
	Vitamin K	0	0
	Bicarbonate	0	0
	Platelets	0	0
	Gastric lavage	0	0

NAC = N-acetyl cysteine; FFP = Fresh frozen plasma.

Table 2: Mean, standard deviation, minimum and maximum of laboratory findings.

Blood parameter	Minimum	Maximum	Mean	SD
ALP	2.50	1056.00	265.04	206.43
Na	134.00	152.00	143.15	3.31
K	.80	9.90	4.07	.86
BUN	.60	76.00	30.72	10.95
Cr	.40	3.70	.91	.38
AST	8.00	5023.00	142.27	601.19
ALT	4.00	4425.00	179.88	704.63
PT	11.00	60.00	14.83	7.43
PTT	23.00	233.00	36.27	25.79
INR	.80	12.50	1.54	1.87

ALP = Alkaline phosphatase; Na = Sodium; K = Potassium; BUN = Blood urea nitrogen; Cr = Creatinine; AST = Aspartate aminotransferase; ALT = Alanine aminotransferase; PT = Prothrombin time; PTT = Partial thromboplastin time; INR = International normalized ratio; SD = Standard deviation.

Table 3: Frequency and percentage of the clinical symptoms in the study subjects.

Symptom	Frequency	%
Nausea, vomiting	101	81.5
Sweating	51	41.1
Abdominal pain	46	37.1
Diarrhea	42	33.9
Blurred vision	37	29.8
Drooling	20	16.1
Fever, shivering	19	15.3
Dizziness	14	11.3
Headache	13	10.5
Weakness, lethargy	7	5.6
Hepatitis	3	2.4
Shortness of breath	3	2.4
Lacrimation	3	2.4
High blood pressure	1	0.8
Low blood pressure	0	0.0
Loss of appetite	1	0.8

Table 4: The laboratory findings in the study subject.

Blood Parameter	Low		Normal		High	
	Frequency	%	Frequency	%	Frequency	%
ALP	0	0	88	71.0	23	18.5
Na	1	.8	88	71.0	30	24.2
K	9	7.3	106	85.5	3	2.4
BUN	3	2.4	101	81.5	13	10.5
Cr	14	11.3	95	76.6	8	6.5
AST	0	0	85	68.5	26	21.0
ALT	0	0	75	60.5	36	29.0
PT	0	0	75	60.5	12	9.7
PTT	0	0	73	58.9	12	9.7
INR	0	0	58	46.8	29	23.4

Normal Ranges: ALP (less than 300); ALT (<300); AST (<300); Na (145-135); K (3.5-5); BUN (15-40); Cr (0.6-1.3); PT (11-15); PTT (20-40); and INR (0.7-1.1).

Table 5: Distribution of the clinical symptoms versus gender among the study subjects.

Symptom	Male		Female		Total	
	Frequency	%	Frequency	%	Frequency	%
Weakness, lethargy	4	5.5	3	5.9	7	5.6
Diarrhea	22	30.1	20	39.2	42	33.9
Nausea and vomiting	59	80.8	42	82.4	101	81.5
Sweating	35	47.9	16	31.4	51	41.1
Blurred vision	17	23.3	20	39.2	37	29.8
Drooling	14	19.2	6	11.8	20	16.1
Low blood pressure	0	0	0	0	0	0
High blood pressure	1	1.4	0	0	1	0.8
Abdominal pain	24	32.9	22	43.1	46	37.1
Headache	10	13.7	3	5.9	13	10.5
Dizziness	7	9.6	7	13.7	14	11.3
Hepatitis	2	2.7	1	2.0	3	2.4
Fever and shivering	12	16.4	7	13.7	19	15.3
Loss of appetite	1	1.4	0	0	1	0.8
Shortness of breath	2	2.7	1	2.0	3	2.4
Lacrimation	1	1.4	2	3.9	3	2.4

Table 6: Distribution of the clinical symptoms in the patients versus their occupations.

Symptom	Self-employed		Retired		Unemployed		Housewife		Student		Univ. student		Worker		Employee		Farmer	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Lethargy	2	5.9	0	0	0	0	3	6.7	0	0	0	0	0	0	1	14.3	1	33.3
Diarrhea	11	32.4	2	5	9	42.9	15	33.3	3	50	0	0	0	0	1	14.3	1	33.3
Nausea, vomiting	28	82.4	4	100	18	85.7	38	84.4	5	83.3	0	0	2	100	3	42.9	3	100
Sweating	18	52.9	2	50	2	9.5	18	40	0	0	2	100	2	100	6	85.7	1	33.3
Blrd. vision	8	23.5	0	0	5	23.8	16	35.6	2	33.3	2	100	1	50	2	28.6	1	33.3
Drooling	7	20.6	0	0	2	9.5	7	15.6	0	0	1	50	1	50	2	28.6	0	0
Low BP	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
High BP	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	33.3
Belly ache	10	29.4	1	25	10	47.6	20	44.4	3	50	0	0	0	0	1	14.3	1	33.3
Headache	5	14.7	1	25	0	0	4	8.9	1	16.7	0	0	0	0	0	0	2	66.7
Dizziness	3	8.8	0	0	0	0	7	15.6	1	16.7	1	50	0	0	1	14.3	1	33.3
Hepatitis	1	2.9	0	0	1	4.8	1	2.2	0	0	0	0	0	0	0	0	0	0
Fever, shiver	7	20.6	0	0	2	9.5	7	15.6	0	0	1	50	1	50	1	14.3	0	0
LOA	0	0	0	0	0	0	0	0	1	16.7	0	0	0	0	0	0	0	0
SOB	1	2.9	0	0	0	0	1	2.2	0	0	0	0	0	0	0	0	1	33.3
Lacrimation	1	2.9	0	0	0	0	2	4.4	0	0	0	0	0	0	0	0	0	0

Blrd. = Blurred; BP = Blood pressure; LOA = Loss of appetite; SOB = Shortness of breath.

Table 7: Distribution of clinical symptoms versus season of poisoning.

Symptom	Spring		Summer		Autumn		Winter	
	Frequency	%	Frequency	%	Frequency	%	Frequency	%
Weakness, lethargy	5	4.4	2	33.3	0	0	0	0
Diarrhea	39	34.5	2	33.3	0	0	1	33.3
Nausea, vomiting	91	80.5	6	100	1	50	3	100
Sweating	49	43.4	1	16.7	1	50	0	0
Blrd. vision	37	32.7	0	0	0	0	0	0
Drooling	20	17.7	0	0	0	0	0	0
Low BP	0	0	0	0	0	0	0	0
High BP	1	0.9	0	0	0	0	0	0
Belly ache	39	34.5	4	66.7	2	100	1	33.3
Headache	12	10.6	1	16.7	0	0	0	0
Dizziness	14	12.4	0	0	0	0	0	0
Hepatitis	3	2.7	0	0	0	0	0	0
Fever/shivering	18	15.9	0	0	0	0	1	33.3
LOA	1	0.9	0	0	0	0	0	0
SOB	3	2.7	0	0	0	0	0	0
Lacrimation	3	2.7	0	0	0	0	0	0

Blrd. = Blurred; BP = Blood pressure; LOA = Loss of appetite; SOB = Shortness of breath

Table 8: Distribution of the clinical symptoms versus underlying diseases.

Symptom	Renal disease		Liver disease		Hematological disease		Diabetes		Hypertension		Coronary disease		Total	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Lethargy	0	0	0	0	0	0	1	14.3	2	18.2	3	37.5	7	5.6
Diarrhea	1	50	0	0	1	50	2	28.6	4	36.4	2	25	42	33.9
Nausea, vomiting	2	100	1	100	2	100	6	85.7	11	100	8	100	101	81.5
Sweating	1	50	0	0	0	0	2	28.6	5	45.5	3	37.5	51	41.1
Blrd. vision	0	0	0	0	0	0	1	14.3	3	27.3	2	25	37	29.8
Drooling	1	50	0	0	0	0	0	0	1	9.1	0	0	20	16.1
Low BP	0	0	0	0	0	0	0	0	0	0	0	0	0	0
High BP	0	0	0	0	1	50	0	0	1	9.1	1	12.5	1	0.8
Belly ache	2	100	0	0	0	0	3	42.9	6	54.5	5	62.5	46	37.1
Headache	0	0	0	0	1	50	0	0	2	18.2	3	37.5	13	10.5
Dizziness	0	0	0	0	0	0	0	0	2	18.2	1	12.5	14	11.3
Hepatitis	0	0	0	0	0	0	0	0	0	0	0	0	3	2.4
Fever/ shivers	0	0	0	0	1	50	0	0	0	0	1	12.5	19	15.3
LOA	0	0	0	0	0	0	0	0	0	0	0	0	1	0.8
SOB	1	50	0	0	1	50	0	0	1	9.1	1	12.5	3	2.4
Lacrimation	0	0	0	0	0	0	1	14.3	1	9.1	0	0	3	2.4

Blrd. = Blurred; BP = Blood pressure; LOA = Loss of appetite; SOB = Shortness of breath.

Table 9: Distribution of the clinical symptoms versus the outcomes.

Symptom	Admission at ward		Consented discharge		ICU admission		Discharge from ER		Death		Referral to specialists		Total	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Lethargy	3	4	2	5.6	1	11.1	0	0	1	100	0	0	7	5.6
Diarrhea	21	28	11	30.6	6	66.7	2	100	1	100	1	100	42	33.9
Nausea, vomiting	61	81.3	28	77.8	9	100	1	50	1	100	1	100	101	81.5
Sweating	31	41.3	18	50	1	11.1	0	0	1	100	0	0	51	41.4
Blrd vision	22	29.3	13	36.1	0	0	2	100	0	0	0	0	37	29.8
Drooling	12	16	8	22.2	0	0	0	0	0	0	0	0	20	16.1
Low BP	0	0	0	0	0	0	0	0	0	0	0	0	0	0
High BP	1	1.3	0	0	0	0	0	0	0	0	0	0	1	0.8
Belly ache	26	34.7	12	33.3	6	66.7	1	50	0	0	1	100	46	37.1
Headaches	8	10.7	4	11.1	1	11.1	0	0	0	0	0	0	13	10.5
Dizziness	11	14.7	3	8.3	0	0	0	0	0	0	0	0	14	11.3
Hepatitis	0	0	0	0	2	22.2	0	0	1	100	0	0	3	2.4
Fever/shiver	13	17.3	4	11.1	1	11.1	0	0	1	100	0	0	19	15.3
LOA	0	0	0	0	0	0	0	0	0	0	1	100	1	0.8
SOB	3	4	0	0	0	0	0	0	0	0	0	0	3	2.4
Lacrimation	2	2.7	1	2.8	0	0	0	0	0	0	0	0	3	2.4

Blrd = Blurred; BP = Blood pressure; ER = Emergency Room; LOA = Loss of appetite; N = Number; SOB = Shortness of breath.