

Research Paper:

Antioxidant and Inhibitory Activities of Enzymes Linked to Type II Diabetes Mellitus: The Novel Role of Chrysobalanus Orbicularis Leaf Extract



Lisa Ilobekemen Ekakitie¹, Oluwafemi Adeleke Ojo^{2*}, Babatunji Emmanuel Oyinloye^{1,3}, Basiru Olaitan Ajiboye¹

1. Phytomedicine, Biochemical Toxicology and Biotechnology Research Laboratories, Department of Biochemistry, Afe Babalola University, Ado-Ekiti, Ekiti State, Nigeria.

2. Department of Biochemistry, Landmark University, Omu-Aran, Kwara State, Nigeria.

3. Biotechnology and Structural Biology (BSB) Group, Department of Biochemistry and Microbiology, University of Zululand, Kwa Dangezwa 3886, South Africa.



How to cite this paper Ilobekemen Ekakitie L, Adeleke Ojo O, Emmanuel Oyinloye B, Olaitan Ajiboye B. Antioxidant and Inhibitory Activities of Enzymes Linked to Type II Diabetes Mellitus: The Novel Role of Chrysobalanus Orbicularis Leaf Extract. Iranian Journal of Toxicology. 2020; 14(3):179-186. <http://dx.doi.org/10.32598/ijt.14.3.371.2>

<http://dx.doi.org/10.32598/ijt.14.3.371.2>



Article info:

Received: 25 Apr 2020

Accepted: 21 Jun 2020

Online Published: 01 Jul 2020

* Corresponding author:

Oluwafemi Adeleke Ojo, PhD.

Address: Department of Biochemistry,

Landmark University, Omu-Aran,

Kwara State, Nigeria.

E-mail: oluwafemiadeleke08@gmail.com

ABSTRACT

Background: Chrysobalanus orbicularis is commonly used as food, the seeds as a spice for the popular pepper soup, and the leaves are traditionally utilized for managing Type II Diabetes Mellitus (T2DM) in the Niger Delta area of Nigeria. Due to the limited scientific evidence on C. orbicularis, this study investigated its antioxidant and inhibitory properties against major enzymes linked to T2DM.

Methods: The antioxidant activity was measured via methods for possible scavenging potentials. The α -amylase and α -glucosidase inhibitory activities were examined using a standard model.

Results: The C. orbicularis aqueous leaf extract compared well with the standard compounds, revealing the high antioxidant and inhibitory properties against α -amylase and α -glucosidase.

Conclusion: The findings validate the conventional application of C. orbicularis in the treatment of patients with type II diabetes mellitus and other oxidative stress-related disorders.

Keywords: Antioxidant activity; C. orbicularis; α -amylase and α -glucosidase; Diabetes mellitus; Medicinal plants

Introduction

Recently, there has been a significant growth in the use of plant-based phytochemicals in the management of diseases and are being propagated in various countries due to the natural origins and low side effects [1]. During oxidative stress, the human body produces free radicals that overwhelm the enzymatic antioxidants, such as Glutathione (GPx) [2]. The

free radicals are mainly produced through biophysical and environmental processes, which cause cellular damages and in most cases result in apoptosis [3]. The quest for natural antioxidants as beneficial agents to inhibit free radicals in the pathogenesis of some human diseases is an important area of clinical interest [4]. Reactive Oxygen Species (ROS) generates hyperglycemia, leading to inequity in antioxidants in the body and results in oxidative stress. However, the blood sugar level is checked by digestive enzymes, like the α -amylase

and α -glucosidase. The enzyme, α -amylase, is responsible for breaking down long-chain carbohydrates while α -glucosidase converts carbohydrate to glucose in the small intestine. The inhibitory effect of α -glucosidase has been acknowledged as a beneficial target for the control of postprandial hyperglycemia and type II diabetes. Therefore, it is of clinical benefit to provide antioxidants, α -amylase and α -glucosidase inhibitors through nutrients that can help manage type II diabetes [5].

Chrysobalanus orbicularis (*C. orbicularis*) is an evergreen plant with a red apple-like fruit locally known as Omillo. The leaves are oval in shape, and the plant belongs to *Chrysobalanaceae* family. It grows mainly as shrub and is known for its management of diabetes mellitus traditionally but has not been fully studied. The whole plant is used locally; its dried fruits and seeds are used for preparing pepper soup, a delicacy in the Southern Nigeria and western Africa. Based on ethno-botanical usage, this study aimed to examine the antioxidant and inhibitory potentials of *C. orbicularis* aqueous leaf extract against α -amylase and α -glucosidase.

Materials and Methods

Plant materials: The leaves from *C. orbicularis* plant were collected from Coco, Itsekiri area in Delta State, Nigeria, and were identified at the Forestry and Research Institute, Ibadan, Oyo State, Nigeria, where they were authenticated and given a voucher Herbarium number (FHI:112232).

Preparation of plant materials: Preparation and extraction of the plant materials were achieved by the procedure defined previously by Nwozo, et al. [6]. The plant leaves were washed with clean water and air dried for three weeks and then pulverized into fine powder in a blender. The powdered sample (1000 g) was extracted in aqueous by cold maceration for 72 hours followed by periodic stirring. The material was freeze-dried and the yield of 23.345g was stored for further analysis.

Biochemical analyses: Assay of Ferric Reducing Antioxidant Potential (FRAP): The FRAP property of aqueous extract of *C. orbicularis* leaves was estimated by evaluating its ability to reduce FeCl_3 solution based on the protocol explained by a previous study [7]. The reducing power was expressed as the gallic acid equivalent.

Assay of 1,1-Diphenyl-1-2-Picryl-Hydrazil (DPPH) radical scavenging ability: The DPPH radical scavenging ability was measured via a previously described procedure [8]. One mL of the extracts at varying concentra-

tions (100-560 $\mu\text{g/mL}$) was incubated at 25°C at dark for 30 minutes with 500 μL of 0.3 mM DPPH solution (prepared in methanol). Then the absorbance was read at 517nm against a blank test tube without the sample.

Assay of Nitric Oxide (NO) scavenging activity: The scavenging activity of the extract of *C. orbicularis* leaves against Nitric Oxide (NO) radical was evaluated by the procedure, as previously described [9]. A 250 μL of the extract at various concentrations (100–500 $\mu\text{g/mL}$) was incubated with 250 μL of 10 mM sodium nitroprusside-sodium phosphate buffer (pH 7.4) for 2h at 37°C . Afterward, 250 μL of Griess reagent was added to the reaction solution and the absorbance was read at 546 nm. The percent inhibition of the NO produced was obtained by comparing the absorbance of the extracts (sample) with that of the control in the absence of scavengers.

Determination of Hydrogen Peroxide (H_2O_2) scavenging activity: The hydrogen peroxide scavenging activity of the aqueous extract of *C. orbicularis* leaves was measured via a previously described procedure [10]. The hydrogen peroxide solution (40 mM) was prepared in phosphate buffer pH 7.4 and its concentration was determined by measuring the absorbance at 560nm, using a UV spectrophotometer. A 1.5mL aliquot of the extract was added to the hydrogen peroxide solution and the absorbance was read at 560nm, using a UV spectrophotometer.

Assay of 2, 2-Azino-Bis-3-ethylbenzotiazolin-6-sulfonic acid (ABTS) scavenging activity: The ABTS activity of the extracts was achieved via the protocol described previously [11]. The stock solutions included 7mM ABTS and 2.4 mM potassium persulfate. The working solution was then prepared by mixing the two stock solutions in equal quantities and allowing them to mix for 14h at 25°C in the dark. The mixture was then diluted by adding 1mL ABTS solution to 60mL methanol, to obtain an absorbance of 0.706 ± 0.01 units at 734nm on a spectrophotometer. Fresh ABTS solution was prepared for each assay. The aqueous extract (1.5mL) was allowed to react with 1mL of the ABTS solution and the absorbance was read at 734nm after 7 min, using a spectrophotometer.

Determination of α -glucosidase inhibitory activity: The inhibitory effect of the aqueous extract of *C. orbicularis* leaves against α -glucosidase activity was determined via a procedure as defined previously [12]. A 250 μL of 1.0 U/mL of α -glucosidase, dissolved in phosphate buffer (100 mmol/L, pH 6.8), was incubated with 250 μL of the extract or acarbose as the standard, for 20 min at 37°C . A 250 μL of p-nitrophenyl- α -D-

glucopyranoside (pNPG) solution (5 mmol/L) that was prepared in the same phosphate buffer was added to the reaction mixture and incubated for another 30 min at 37°C. The absorbance of released p-nitrophenol was measured at 405 nm.

Assay of α -amylase inhibitory activity: The inhibitory action of the extract against α -amylase activity was evaluated based on the established procedure of a previous study [13]. Briefly, 250 μ L (1.5 mg/mL stock) of the extracts at varying concentrations (100–500 μ g/mL) or acarbose was incubated with 250 μ L of the enzyme (porcine pancreatic amylase 2 U/mL) in phosphate buffer (100 mmol/L, pH 6.8) for 20 min at 37°C. Thereafter, 250 μ L of 1% starch prepared in the same buffer was added to the premixed solution and incubated further for 1h at 37°C. One mL of the color reagent, Dinitrosalicylate (DNS), was added and the mixture was boiled for 10 min. The absorbance was read at 540nm and the inhibitory activity was calculated as the proportion of sample versus the control.

Data analyses: The data were analyzed via GraphPad Prism 8.0 (Version 8, Software Program, GraphPad Prism Inc., San Diego, CA, USA) and presented as the Mean \pm SD in triplicates.




Figure 1. FRAP scavenging ability of aqueous extract of *C. orbicularis* leaves

Values are shown as Mean \pm SD; n=10;

Legend: AACO - *Chrysobalanus orbicularis* leaf extract; Gallic acid is the standard;

FRAP- Ferric reducing antioxidant potential; a-eSuperscripted alphabets over the bars for a given concentration of each extract indicate significance of difference at P<0.05.

Results

The FRAP scavenging activity of the aqueous extract of *C. orbicularis* leaves is shown in Figure 1. The activity showed a rise with increasing concentration of the extract. In contrast, gallic acid revealed a significant increase in reducing ability as compared to the extract (P<0.05). The DPPH scavenging activity of the extract exhibited inhibitory activity and compared well with butylated hydroxytoluene, used as the standard (Figure 2).

The nitric oxide scavenging activity rose progressively with increasing concentration and compared well with that of the standard (Figure 3). As presented in Figure 4, an upsurge in the concentration of the extract yielded an increase in hydrogen peroxide scavenging activity and compared well with that of the gallic acid as the standard. Figure 5 shows the result of the scavenging activity of ABTS. There was an upsurge with the increasing concentration, and compared well with the standard. Figure 6 shows the result of the inhibitory activities of the extract against α -amylase and α -glucosidase, and compared well with acarbose as the standard.




Figure 2. DPPH scavenging ability of aqueous extract of *C. orbicularis* leaves

Values are represented as Mean \pm SD; n=10;

Legend: AACO - *Chrysobalanus orbicularis* leaf extract; BHT is the standard (butylated hydroxytoluene);

DPPH- 1, 1-diphenyl-2-picryl-hydrazil. a-cSuperscripted alphabets over the bars for a given concentration of each extract indicate significance of difference at P<0.05.




Figure 3. Nitric oxide scavenging ability of aqueous extract of *C. orbicularis* leaves

Values are represented as Mean±SD; n=10;

Legend: AACO = *Chrysobalanus orbicularis* leaf extract. BHT is the standard (butylated hydroxyl toluene);

a-eSuperscripted alphabets over the bars for a given concentration of each extract indicate significance of difference at P<0.05.

Discussion

Free radicals have been involved in the development and progression of numerous diseases, such as diabetes and its complications [14-17]. It has been reported that the hyperglycemia produces free radicals and results in an imbalance of radical-antioxidant system in favor of radicals, leading to oxidative stress [18, 19]. The ability of a constituent to scavenge free radicals is linked to its electron transfer ability [20]. In the present study, the free radical scavenging and ferric reducing properties of the aqueous extract of *C. orbicularis*




Figure 4. H₂O₂ scavenging ability of the aqueous extract of *C. orbicularis* leaves

Values are represented as Mean±SD; n=10;

Legend: AACO – *Chrysobalanus orbicularis* leaf extract. BHT is the standard (butylated hydroxyl toluene);

a-bSuperscripted alphabets over the bars for a given concentration of each extract indicate significance of difference at P<0.05.

leaves were evaluated in vitro. The scavenging property was investigated to test the total antioxidant capacity, using ferric reducing power, 1,1-diphenyl-1-2-picryl-hydrazil (DPPH), hydrogen peroxide (H₂O₂), Nitric Oxide (NO) and 2, 2-Azino-bis-3-ethylbenzotiazolin-6-sulfonic acid (ABTS). These methods are generally employed to assess the efficacy of the antioxidant activities of plants' extracts [8, 13]. The aqueous extract of *C. orbicularis* showed high levels of radical scavenging activities, thus this plant possesses antioxidant potentials. This property is likely to offer inhibitory effect and;




Figure 5. ABTS scavenging ability of the aqueous extract of *C. orbicularis* leaves

Values are represented as Mean±SD; n=10;

Legend: AACO: *Chrysobalanus orbicularis* leaf extract. BHT is the standard (butylated hydroxyl toluene);

a-dSuperscripted alphabets over the bars for a given concentration of each extract indicate significance of difference at p<0.05.




Figure 6. α -amylase and α -glucosidase inhibitory activity of *C. orbicularis* leaf extract

Legend: AACO – *Chrysobalanus orbicularis* leaf extract;

ACARBOSE: The standard values are represented as mean \pm standard deviation (SD); n=10;

a-eSuperscripted alphabets over the bars for a given concentration of each extract indicate significance of difference at P<0.05.

therefore, used clinically in the management of the oxidative stressors in such diseases, as diabetes mellitus type II.

Delaying the postprandial rise in the serum glucose level via the inhibition of α -amylase and α -glucosidase is critical to the management of patients with diabetes type II [21, 22]. Specifically, the inhibition of the enzymes that are implicated in the intermediary metabolism of carbohydrates is one of the most effective therapeutic approach in lowering postprandial blood glucose levels in diabetic patients [23, 24]. The enzyme, α -amylase, has been known to control and hydrolyze complex carbohydrates to disaccharides, while α -glucosidase breaks down oligosaccharides to monosaccharides, thereby leading to a rise in postprandial serum glucose level [25, 26]. The ability of *C. orbicularis* leaf extract to inhibit the two enzymes confirms the potential of the extract in the management of diabetes mellitus [18, 19]. Since the antioxidant activity of the extract has not been examined previously, this study attempted this investigation for the first time. The antioxidant activity of plants is primarily linked to their phenolic compounds content.

In a study by de Oliveira Barbosa, et al. [27], the extract was rich in phenolic compounds with high antioxidant properties. The results obtained by the current study are consistent with those reported by de Oliveira Barbosa, et al. [27]. Further, the high scavenging property has been attributed to the hydroxyl groups present in the chemical structure of the phenolic compounds, which are the essential components for scavenging free radicals [28, 29].

Conclusions

The aqueous extract of *C. orbicularis* leaves exhibited high antioxidant activity in vitro compared to the standard. In addition, the extract showed inhibitory property against α -amylase and α -glucosidase. These findings suggest that the extract of *C. orbicularis* leaves has the ability to inhibit the oxidative stress induced by free radicals in various human pathologic conditions, such as diabetes mellitus. This study is the first to report the in vitro antioxidant activity of the extract against α -amylase and α -glucosidase.

Ethical Considerations

Compliance with ethical guidelines

The study is an in vitro one which did not involve animals and human subjects. All ethical principles of chemicals and plants studies were applied in this research.

Funding

This study was funded by the authors, without receiving any financial support from private or public organizations.

Author's contributions

Conceptualization and methodology: Basiru Olaitan Ajiboye, Oluwafemi Adeleke Ojo; Data analysis: Lisa Ilobekemen Ekakitie; Writing of the original draft, review and ed-

iting of the manuscript: Lisa Ilobekemen Ekakitie, Basiru Olaitan Ajiboye, Babatunji Emmanuel Oyinloye.

Conflict of interest

The authors declare no conflict of interests with any entity in conducting this study.

Acknowledgements

The authors wish to acknowledge the support of the staff and management of the Departments of Biochemistry at Afe Babalola and Landmark Universities, in Nigeria, and the Department of Biochemistry and Microbiology at University of Zululand in South Africa.

References

- [1] Kooti W, Farokhipour M, Asadzadeh Z, Ashtary-Larky D, Asadi-Samani M. The role of medicinal plants in the treatment of diabetes: A systematic review. *Electron Physician*. 2016; 8(1):1832-42. [DOI:10.19082/1832] [PMID] [PMCID]
- [2] Oyinloye BE, Adenowo AF, Kappo AP. Reactive oxygen species, apoptosis, antimicrobial peptides and human inflammatory diseases. *Pharmaceuticals*. 2015; 8(2):151-75. [DOI:10.3390/ph8020151] [PMID] [PMCID]
- [3] Ibitoye OB, Ghali UM, Adekunle JB, Uwazie JN, Ajiboye TO. Antidyslipidemic, anti-inflammatory, and antioxidant activities of aqueous leaf extract of *Dioscorea oppositifolia* (Stapf) Diels in high-fat diet-fed rats. *Evid Based Complement Alternat Med*. 2017; 2017:8128125 [DOI:10.1155/2017/8128125] [PMID] [PMCID]
- [4] Idamokoro EM, Masika PJ, Muchenje V. A report on the in vitro antioxidant properties of *Vachellia karroo* leaf extract: A plant widely grazed by goats in the Central Eastern Cape of South Africa. *Sustainability*. 2017; 9(2):164. [DOI:10.3390/su9020164]
- [5] Garvey WT. Phentermine and topiramate extended-release: A new treatment for obesity and its role in a complications-centric approach to obesity medical management. *Expert Opin Drug Saf*. 2013; 12(5):741-56. [DOI:10.1517/14740338.2013.806481] [PMID] [PMCID]
- [6] Nwozo SO, Oyinloye BE. Hepatoprotective effect of aqueous extract of *Aframomum metegueta* on ethanol-induced toxicity in rats. *Acta Biochim Pol*. 2011; 58(3):355-8. [DOI:10.18388/abp.2011_2246] [PMID]
- [7] Pulido R, Bravo L, Saura-Calixto F. Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/antioxidant power assay. *J Agric Food Chem*. 2000; 48(8):3396-402. [DOI:10.1021/jf9913458] [PMID]
- [8] Ojo O, Oloyede O, Olarewaju O, Ojo A, Ajiboye B, Onikanni S. In-Vitro Antioxidant and Free Radical Scavenging Activities of *Ocimum gratissimum*. *World J Pharm Res*. 2013; 2(6):1899-912. <https://www.researchgate.net/publication/258240873>
- [9] Marcocci L, Maguire JJ, Droylefaix MT, Packer L. The nitric oxide-scavenging properties of Ginkgo biloba extract EGB 761. *Biochem Biophys Res Commun*. 1994; 201(2):748-55. [DOI:10.1006/bbrc.1994.1764] [PMID]
- [10] Ruch RJ, Cheng SJ, Klaunig JE. Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea. *Carcinogenesis*. 1989; 10(6):1003-8. [DOI:10.1093/carcin/10.6.1003] [PMID]
- [11] Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med*. 1999; 26(9-10):1231-7. [DOI:10.1016/S0891-5849(98)00315-3]
- [12] Ademiluyi AO, Oboh G. Soybean phenolic-rich extracts inhibit key-enzymes linked to type 2 diabetes (α -amylase and α -glucosidase) and hypertension (angiotensin I converting enzyme) in vitro. *Exp Toxicol Pathol*. 2013; 65(3):305-9. [DOI:10.1016/j.etp.2011.09.005] [PMID]
- [13] Shai LJ, Masoko P, Mokgotho MP, Magano SR, Mogale AM, Boaduo N, et al. Yeast alpha glucosidase inhibitory and antioxidant activities of six medicinal plants collected in Phalaborwa, South Africa. *S Afr J Bot*. 2010; 76(3):465-70. [DOI:10.1016/j.sajb.2010.03.002]
- [14] Ojo OA, Ajiboye B, Fadaka A, Taro P, Shariati MA. Nrf2-Keap1 activation, a promising strategy in the prevention of cancer. *Free Radic and Antioxid*. 2017; 7(1):01-07. [DOI:10.5530/fra.2017.1.1]
- [15] Phaniendra A, Jestadi DB, Periyasamy L. Free radicals: Properties, sources, targets, and their implication in various diseases. *Indian J Clin Biochem*. 2015; 30(1):11-26. [DOI:10.1007/s12291-014-0446-0] [PMID] [PMCID]
- [16] Kehrer JP, Klotz LO. Free radicals and related reactive species as mediators of tissue injury and disease: Implications for health. *Crit Rev Toxicol*. 2015; 45(9):765-98. [DOI:10.3109/10408444.2015.1074159] [PMID]
- [17] Ojo OA, Afon AA, Ojo AB, Ajiboye BO, Oyinloye BE, Kappo AP. Inhibitory Effects of Solvent-partitioned fractions of two Nigerian herbs (*Spondias mombin* Linn. and *Mangifera indica* L.) on α -amylase and α -glucosidase. *Antioxidants*. 2018; 7(6):73. [DOI:10.3390/antiox7060073] [PMID] [PMCID]
- [18] Palsamy P, Subramanian S. Ameliorative potential of resveratrol on proinflammatory cytokines, hyperglycemia mediated oxidative stress, and pancreatic β -cell dysfunction in streptozotocin-nicotinamide-induced diabetic rats. *J Cell Physiol*. 2010; 224(2):423-32. [DOI:10.1002/jcp.22138] [PMID]
- [19] Ojo OA, Ajiboye BO, Imiere OD, Adeyolu O, Olayide I, Fadaka A. Antioxidative properties of *Blighia sapida* K.D. Koenig stem bark extract and inhibitory effects on carbohydrate hydrolyzing enzymes associated with non-insulin dependent diabetes mellitus. *Pharmacogn J*. 2018; 10(2):376-83. [DOI:10.5530/pj.2018.2.63]
- [20] Ojo OA, Ojo AB, Ajiboye BO, Olaiya O, Akawa A, Olaoye O, et al. Inhibitory effect of *Bryophyllum pinnatum* (Lam.) Oken leaf extract and their fractions on α -amylase, α -glucosidase and cholinesterase enzyme. *Pharmacogn J*. 2018; 10(3):497-506. [DOI:10.5530/pj.2018.3.82]
- [21] Ajiboye BO, Ojo OA, Oyinloye BE, Akuboh O, Okesola MA, Idowu O, et al. In vitro antioxidant and inhibitory activities of polyphenolic-rich extracts of *Syzygium cumini* (Linn)

- Skeels leaf on two important enzymes relevant to type II diabetes mellitus. *Pak J Pharm Sci.* 2020; 33(2):523-9. [PMID]
- [22] Ajiboye BO, Ojo OA, Okesola MA, Oyinloye BE. Ethyl acetate leaf fraction of *Cnidioscolus aconitifolius* (Mill.) I. M. Johnst: Antioxidant potential, inhibitory activities of key enzymes on carbohydrate metabolism, cholinergic, monoaminergic, purinergic, and chemical fingerprinting. *Int J Food Prop.* 2018; 21(1):1697-715. [DOI:10.1080/10942912.2018.1504787]
- [23] Ilahi I, Samar S, Khan I, Ahmad I. In vitro antioxidant activities of four medicinal plants on the basis of DPPH free radical scavenging. *Pak J Pharm Sci.* 2013; 26(5):949-52. [PMID]
- [24] Ajiboye BO, Ojo OA, Adeyonu O, Imiere O, Olayide I, Fadaka A, et al. Inhibitory effect of key enzymes relevant to acute type-2 diabetes and antioxidative activity of ethanolic extract of *Artocarpus heterophyllus* stem bark. *J Acute Dis.* 2016; 5(5):423-9. [DOI:10.1016/j.joad.2016.08.011]
- [25] Ojo OA, Ojo AB, Ajiboye B, Olayide I, Fadaka A. *Helianthus annuus* leaf ameliorates postprandial hyperglycaemia by inhibiting carbohydrate hydrolyzing enzymes associated with type-2 diabetes. *Iran J Toxicol.* 2016; 10(5):17-22. [DOI:10.29252/arakmu.10.5.17]
- [26] Ajiboye BO, Ojo OA, Okesola MA, Akinyemi AJ, Talabi JY, Idowu OT, et al. In vitro antioxidant activities and inhibitory effects of phenolic extract of *Senecio biafrae* (Oliv and Hiern) against key enzymes linked with type II diabetes mellitus and Alzheimer's disease. *Food Sci Nutr.* 2018; 6(7):1803-10. [DOI:10.1002/fsn3.749] [PMID] [PMCID]
- [27] de Oliveira Barbosa AP, de Oliveira Silveira G, de Menezes IAC, Neto JMR, Bitencurt JLC, dos Santos Estavam C, et al. Antidiabetic effect of the *Chrysobalanus icaco* L. aqueous extract in rats. *J Med Food.* 2013; 16(6):538-43. [DOI:10.1089/jmf.2012.0084] [PMID]
- [28] Nahak G, Sahu RK. Antioxidant activity in bark and roots of neem (*Azadirachta indica*) and mahaneem (*Melia azedarach*). *Cont J Pharm Sci.* 2010; 4:28-34. <https://www.researchgate.net/publication/215613224>
- [29] Ajiboye BO, Ojo OA, Fatoba B, Afolabi OB, Olayide I, Okesola MA, et al. In vitro antioxidant and enzyme inhibitory properties of n-butanol fraction of *Senna podocarpa* (Guill. and Perr.) leaf. *J Basic Clin Physiol Pharmacol.* 2020; 31(1):20190123. [DOI:10.1515/jbcpp-2019-0123] [PMID]

This Page Intentionally Left Blank