

Research Paper: Assessment of Non-carcinogenic Risk of Nitrate in Agricultural Products



Abbas Alimoradian¹, Reza Tajik², Mohammad Jamalian³, Mohaddeseh Asafari^{4*}, Rahmatolah Moradzadeh⁵

1. Department of Pharmacology, School of Medicine, Arak University of Medical Sciences, Arak, Iran.
2. Department of Occupational Health Engineering, School of Public Health, Arak University of Medical Sciences, Arak, Iran.
3. Department of Forensic Medicine, School of Medicine, Arak University of Medical Sciences, Arak, Iran.
4. Deputy of Food and Drug, Arak University of Medical Sciences, Arak, Iran.
5. Department of Epidemiology, School of Health, Arak University of Medical Sciences, Arak, Iran.



How to cite this paper Alimoradian A, Tajik R, Jamalian M, Asafari M, Moradzadeh R. Assessment of Non-carcinogenic Risk of Nitrate in Agricultural Products. Iranian Journal of Toxicology. 2021; 15(4):257-264. <http://dx.doi.org/10.32598/ijt.15.4.784.2>

doi <http://dx.doi.org/10.32598/ijt.15.4.784.2>



Article info:

Received: 24 Apr 2021

Accepted: 19 Jun 2021

Online Published: 01 Oct 2021

* Corresponding author:

Mohaddeseh Asafari, MSc.

Address: Deputy of Food and Drug, Arak University of Medical Sciences, Arak, Iran.

E-mail: m.asafari@arak.ac.ir

ABSTRACT

Background: Nitrate content is one of the most critical factors to determine the quality of vegetables, and its permissible limits in food chain is important to the human health. Due to the harmful effects of nitrate, many studies have been conducted on its accumulation in crops in recent years. This study assessed the nitrate concentrations in some edible vegetables and the associated health risks.

Methods: In this descriptive study, raw agricultural produce, such as leafy vegetables, tubers, cucurbits, kitchen produce and edible grains were collected in the winter and summer. The samples were then transferred to a laboratory by maintaining the temperature-controlled supply chain. After preparing the samples, the nitrate content was determined in each, using an Ultraviolet-Visible spectrophotometric (UV/Vis) unit. The data were analyzed statistically on SPSS v. 26.

Results: The mean nitrate contents in leafy and kitchen produce, grains, and tuber vegetables were approximately 130, 48, 101, and 61 ppm, respectively. The average nitrate content in the winter was around 38 ppm and in the summer about 44 ppm. The highest nitrate content was documented in spinach (1100.15 ppm) and the lowest in tomatoes (20.97 ppm).

Conclusion: The results indicated that the highest nitrate content was found in leafy produce grown in northern Iran. The highest health risk for non-carcinogenic conditions was likely to be linked to the consumption of spinach and other edible vegetables, wheat, rice, and potatoes.

Keywords: Dietary intake, Fruits, Vegetables, Health risk assessment, Nitrate content

Introduction

Nitrate content in vegetables and edible crops is one of the harmful chemicals for human health [1]. Nitrate and nitrite are two anions in the nitrogen cycle that are widely distributed in nature. Due to the association of nitrate with methemoglobinemia, a fatal disease, the consumption of foods and beverages

with high nitrate contents (0.3% by weight) is harmful to human health, and is linked to certain types of gastrointestinal cancers [2, 3]. Nitrite may bind food amines to form N-nitrosamine, associated with various types of cancers, including esophageal and stomach neoplasia [3, 4]. According to studies, all edible vegetables contain nitrates in varying amounts, ranging from 1 to 10,000 mg/kg [4, 5]. Depending on the type of vegetables, the nitrate contamination varies based on the nitrogen con-

tent in the soil, temperature, sunlight, and other factors [5]. Edible vegetables, such as potatoes, are among the primary sources of dietary nitrate-containing foods and account for 85% of nitrate intake in humans [4, 6, 7].

Nitrate content is one of the most important factors in determining the quality of vegetables. Nowadays, due to excessive use of nitrogen-containing fertilizers to enhance the vegetations' growth, many vegetables contribute a high percentage of nitrate to the human diet [8]. Humans receive the majority of their nitrate by consuming contaminated food products and vegetables. Unfortunately, in many parts of Iran, cucumbers, potatoes, and watermelons have been reported to have high nitrate contents [9]. Considering that edible vegetables and other kitchen produce absorb and retain large amounts of nitrite and nitrate, the consumption of such products by humans can endanger their health [9]. Nitrate is a major pollutant that accumulates in food chain mainly due to the use of imbalanced nitrogenous fertilizers in farming industry.

Nitrate is converted to nitrite in the digestive system of humans and animals, by binding to some amino acids to produce nitrosamine, which is a carcinogenic chemical [10]. The International Agency for Research on Cancer (IARC) has classified nitrite in foods as a carcinogen of group 2A [11]. This agency determines the permissible limits of the hazardous substances in agricultural products and the environment [11]. The Admissible Daily Intake (ADI) of nitrate for humans through food and water consumption is between zero and 3.7 mg/kg of body weight, as set by the World Health Organization and the European Union Scientific Committee [12]. Accordingly, a person with an average weight of 70 kg should not consume more than 259 mg nitrate per day.

Fertilizers used in agriculture are the most important causes of nitrate contamination. The presence of nitrate in agricultural materials causes the associated environmental and human health problems [13]. Due to its adverse effects on human health, much attention has been paid to the accumulation of this chemical compound in edible vegetables and garden produce. Nitrogen is absorbed by plants in two chemical forms: ammonium and nitrate; however, plants prefer adsorbing nitrate compounds [14]. The nitrate content in plants is influenced by environmental and genetic factors [15]. Nitrate is non-toxic inherently and is essential in human and animal diets. However, its metabolites and byproducts, such as nitrite, nitric oxide, and nitrous compounds, can have harmful effects on human health [16]. Due to the contamination of soil and agricultural products with nitrous compounds, the contents of nitrate and nitrite in vegeta-

bles have become one of the public health concerns in recent years. The objective of this study was to assess the nitrate contents in popular agricultural products, both at the supplier level and in general consumption by people.

Materials and Methods

Sampling: To collect the information on nitrate contents in 18 agricultural products, we contacted the main greengrocers in Arak, and completed a detailed form based on the available documents. The products were leafy fruits and vegetables, tuber, kitchen produce, grains and cereals. These included leafy vegetables, lettuce, spinach, cabbage, potatoes, carrots, onions, sugar beets, beetroots, eggplants, watermelons, melons, tomatoes, edible mushrooms, pinto beans, soy protein, wheat and rice were sampled each month if available. The sample size was determined based on the cargo bills, and the suppliers information in the summer and winter of 2019. The sampling occurred at least 9 times per season, collecting about 1-3 kilog of each product, containing at least 10 pieces of each product per kilogram [17]. The samples were transferred to Marjaan Khatam Laboratory in Tehran on a monthly basis, and stored in a cold room at optimal conditions, with each group bearing its identification tag and number.

Sample preparation & laboratory Methods: The determination of the nitrate levels were carried out according to established standards, using a uv/vis spectrophotometer (Shimatdzu; Japan). Initially, the edible parts of the produce were separated and washed with regular and then deionized water to prepare the study samples. The samples were dried, crushed, and homogenized, and a 100g of each sample was collected and placed in an incubator at 80°C, with the drying time dependant on the desired texture of the sample. For example, thin vegetables (lettuce, celery, cabbage, and radish leaves) needed 12 hours, while vegetables with thick and fleshy texture (potatoes, tomatoes, and radish tubers) took 24 hours to dry completely. In the case of watermelons and cantaloups, their extracts were prepared, using a grinder and then measured. Ten grams of these produce, such as cantaloupe, was added to 25 mL 2% acetic acid, shaken for 30 minutes on a rotary shaker then centrifuged for three minutes at 5000 rpm, filtered through Wattman's paper, and read at 400 nm on a uv/vis spectrophotometer. The experiments followed the instructions provided by the Institute of Standards and Industrial Research of Iran (document No. 4106) [12].

Statistical analysis: Due to the abnormal distribution of nitrate in the samples, nonparametric statistical tests




Figure 1. Mean nitrate contents in products compared with the allowable limits as set by ISIRI.

were used for the analyses. For each product, the nitrate content was checked and compared for winter versus summer, using the Mann-Whitney test. For comparison among the samples, we used Kruskal-Wallis test. To analyze the data, we grouped them by the product's region and the type. The farming regions were divided into five central areas (Alborz, Dezful, Isfahan, Hamedan, Markazi, Qazvin), West (Khorramabad, Ahvaz), East (Bojnourd, Mashhad), North (Mazandaran), and South (Bandar Abbas, Shiraz, Chabahar). The crops were categorized into leafy (lettuce, vegetables, spinach & cabbage), kitchen produce (watermelon, melon, tomato, eggplant & edible mushroom), grains (beans, soy beans, rice & wheat), tuber vegetables (onions, potatoes, carrots, sugar beets & root beets).

Risk assessment: To assess the human health risk of non-cancerous diseases, we used the formula provided by the U.S. Environmental Protection Agency (USEPA) [17-19]. First, we calculated the number of contaminants absorbed in the products per kilogram of body weight per day, using the following Equations:

$$EDI = (CF \times IR \times EF \times ED) / (BW \times AT)$$

$$EDI = \text{Daily absorption of pollutant}$$

$$CF = \text{Concentration of contaminants in food}$$

$$IR = \text{Daily consumption}$$

$$FI = \text{amount of pollutants absorbed by the body through food}$$

This coefficient varied between 0.25 and 0.4 [17].

$EF = \text{the frequency of consumption per year}$

$ED = \text{the number of years used this food}$

In this study was 33.2.

$BW = \text{Body weight}$

In this study was considered 55.9 kg for adults and 15 kg for children [20].

$AT = \text{Multiply the ED by the number of days per year}$

$THQ = EDI / RFD_i$

Was used to determine the risk of non-cancerous diseases [17].

$RFD = \text{maximum tolerance}$

The following Equation was also used to assess the extent of exposure [21]:

$$\text{Exposure assessment} = \text{Concentration of the hazard (ng/g)} \times \text{Food consumption (g/day/w)}$$

Results

Due to the normal distribution of the data derived from Kolmogorov-Smirnov test, we used nonparametric tests for the statistical analyses. The average amount of nitrate measured in the products was 165 ± 27.88 ppm.

Table 1. The nitrate contents in various products versus seasons and locations

Variables	Descriptive Statistics				
	N	Mean	λ^*	P	
Nitrate per Plants	Leafy Vegetables	21	129.74	71.75	0.0001
	Kitchen Produce	51	47.95		
	Legumes and Cereals	18	101.25		
	Root Vegetables	52	60.78		
Nitrate per Location	Center	118	70.83	9.137	0.058
	South	8	60.38		
	North	9	106.78		
	East	3	53.17		
	West	4	48.00		
Nitrate per Season	Warm season	38	43.45	686	0.281
	Cold season	42	37.83		

* λ =Kruskal-Wallis test.

The means for nitrate contents in the products and their permissible limits, based on the national standard, are shown in [Figure 1](#).

The results of nitrate contents in leafy, tuber, root, and kitchen produce samples for both summer and winter are shown in [Table 1](#). As evident by this table, the highest contamination of nitrate was related to leafy vegetables, followed by the grains and cereal samples. Geographically, the highest nitrate content was from the northern regions and the lowest found in the western areas of Iran. Also, the nitrate contents of the crops were lower for the warm season than that of the cold season.

Assessment of exposure and undesirable effects:

Based on the results of the exposure assessment, the high consumptions were for tuber vegetables (19500), lettuce (18000), tomato (13080) (g/day/w), and the low consumptions belonged to mushrooms (42), legumes (1900), onion (3510) (g/day/w) and the average consumption is 1910.3 (g/day/w).

Pathogenic rates: The potential pathogenic risks of the products are presented in [Table 2](#). According to the formula for Target Hazard Quotient (THQ), most of the products carried their risk of pathogenicity. Also, the nitrate aggregation was higher than the permissible limits in such products as leafy vegetables, rice, wheat, and potatoes. However, due to the cumulative effects of ni-

trate over time and consumption of other foods, such as ready-made meat products [22] and drinking water [23], there is also a risk of non-cancerous pathogenicity for other products, such as melon, carrot and eggplant.

Discussion

In this study, nitrate exposure was assessed for food safety, based on the average and highest concentrations of the contaminant in different regions. The results showed that the mean exposure rate in the northern areas was the highest (106.78), with the western regions having the lowest rate (40) of nitrate contamination ($P=0.05$). This suggests that the soil type, fertilization practices, rainfall and irrigation, cultivation frequency and type, differences in geographic locations, absorption rate, and using chemical fertilizers influence the nitrate accumulation in the vegetables. According to other reports, nitrate from fertilizers quickly penetrates into light and permeable soils. This reduces the nitrate contents in plants. However in heavy and less permeable soils, nitrate remains in the plant roots, and the content in them rises. For this reason, in farms with light soils, less fertilizer remains available and accumulate in crops [24, 25]. The most contaminated samples with nitrate were leafy products (129.74 ppm) with the least contaminated ones (47.95 ppm) being the tuber vegetables. This suggests that nitrite and nitrate contents vary with the type of vegetables and products. The important factors include genetic structure,

Table 2. Potential severity of non-carcinogenic risks of the agricultural products

Fruites	Target Hazard Quotient (THQ)	Level of Risk
Spinach	16.6	High risk
Vegetables	13.7	High risk
Wheat	8.1	High risk
Rice	3.4	Medium
Potato	2.6	Medium
Lettuce	2.5	Medium
Melone	1.5	Low risk
Carrot	1.16	Low risk
Eggplant	1.1	Low risk
Watermelone	0.9	No risk
Cabbage	0.9	No risk
Tomato	0.5	No risk
Onion	0.2	No risk

being tuber or leafy, root structure, water absorption rate, and acidity or alkalinity of the product [9].

The other studies have reported that the amount of nitrate in vegetables varies from country to country and in different regions of the same city or country [26-28]. Hasani Moghadam et al. [8] have reported that nitrate levels are higher in tuber parts than that in the plant leaves. Their finding is inconsistent with our results. While Shahbaz-zadegan [29] and Pourmoghimi [30] reported that the order of nitrate contents was highest in leafy vegetables followed by roots and tubers, and the lowest in vegetables. Further, the highest nitrate contamination was observed in the spring, which is inconsistent with those reported earlier by Alemzadeh [31] and Doomari [25], who reported higher nitrate accumulation in plants in the winter. However, many studies elsewhere in the world have reported that temperature negatively affects the plant growth and influences the nitrate content [24, 32-34]. Colla et al. have suggested that crops produced in the late spring contained more nitrate than those in the winter [24].

Exposure to diets contaminated with nitrate and nitrite results in the formation of nitrosoamines, thus becoming a significant public health issue. This is likely to lead to acute and chronic conditions, such as methemoglobinemia, thyroid disorders and some forms of cancer

[26]. Previous studies have shown that nitrate can impair iodine absorption by the thyroid gland, leading to its hypertrophy [35]. Other pathological conditions may also occur, such as gastrointestinal disorders, GI reflux, colon and oral cancers [36, 37]. It has been reported that some other compounds in vegetables, such as ascorbic acid, phenols and tocopherols, carotenoids, and flavonoids can prevent the formation of N-nitro compounds and have the ability to inhibit nitro substance [26]. Based on the findings of the current study, the highest risk of pathogenicity was related to the consumption of contaminated spinach, vegetables, wheat, rice and potatoes. Watermelon, cabbage, tomato and onion were reported to have none to little risk of pathogenicity. Considering the products with the highest risk, wheat, rice, potatoes and leafy vegetables, play an important role in the human food basket. Based on the food basket for each Iranian person per day, 310 g of bread, 95 g of rice, 70 g of potatoes, 300 g of vegetables, and 280 g of fruit are considered the desirable amounts [38].

Recently, Uddin et al. examined the nitrate levels in fruits and vegetables, and assessed the health risks. They have claimed that the pathogenicity risk of almost all samples except for radish was within the acceptable range [39]. One study has reported the risk of THQ less than one [40], which is inconsistent with the results of the current study. Thomson et al. [41] assessed the risk

of nitrate in foods and drinking water, and reported that the nitrate levels in vegetables, e.g., broccoli, exceeded the permissible limit and that more than half of people in New Zealand were at risk of the side effects due to high ADI intakes. In the guidelines of the Iranian National Standards, despite the lower per capita vegetable consumption compared to those in other countries, has reported that the high-risk range of nitrate in all of the popular vegetables consumed by average Iranian household is considered much lower than those of Asian, European and American countries [15]. Based on the maximum daily absorption of nitrate per kilogram of body weight, as set by the World Health Organization [42], consuming more than recommended limits causes nitrate-induced disorders and harmful effects on the body. It is noteworthy that this issue is more critical in children, and the amount is reduced to 62 g per day for a 25 kg child. Also, different parts of vegetables and the plants' age can influence the nitrate level. The nitrate content in young and internal parts has been lower than the old and external parts of many products [40].

Conclusions

This study demonstrated that the leafy vegetables contained high amounts of nitrate, which was more than the maximum limit set by the Iranian National Standards. Although the average amount of nitrate in spinach, vegetables, wheat, rice, and potatoes was permissible, it can be concluded that the risk of pathogenicity increases due to the frequency of their consumption over time. In light of the fact that nitrate is carcinogenic, the human exposure to this compound is worrisome. Considering the high contents of nitrate in some crops being above the permissible limits, the agricultural management and harvesting methods should be monitored, and effective steps taken to reduce nitrate levels in edible produce. Future research should also focus on the effects of nitrate in various populations, such as infants and children, adults over 65 years old, pregnant and lactating women, and people with different health conditions and/or diets.

Ethical Considerations

Compliance with ethical guidelines

The study did not involve human participants or animals, therefore, the ethical guidelines did not apply. The study protocol was approved by Arak University Research Council (Registration #: IR.ARAKMU.REC.1399.270).

Funding

This paper was part of a comprehensive plan to monitor the agricultural products at Arak University of Medical Sciences (No.: IR.ARAKMU.REC.1399.270). This research was supported by Deputy of Food and Drug of Arak University of Medical Science.

Author's contributions

All authors equally contributed to preparing this article.

Conflict of interest

The authors declare no conflict of interests.

Acknowledgments

The authors express their gratitude to the Research Council, Arak University of Medical Sciences for funding this research project. We also appreciate Marjaan Khatam Laboratory in Tehran, Iran, for their assistance with the experiments.

References

- [1] Huang GW, Liu H, Li X, Ma M. Exploring drivers of nitrate contamination of drinking water in an arid region of China. *J Environ Inform.* 2019; 33(2):105-12. [DOI:10.3808/jei.201900409]
- [2] Hmelak Gorenjak A, Cencič A. Nitrate in vegetables and their impact on human health. A review. *Acta Aliment.* 2013; 42(2):158-72. [DOI:10.1556/AAlim.42.2013.2.4]
- [3] Ding Z, Johanningsmeier SD, Price R, Reynolds R, Truong VD, Payton SC, et al. Evaluation of nitrate and nitrite contents in pickled fruit and vegetable products. *Food Control.* 2018; 90:304-11. [DOI:10.1016/j.foodcont.2018.03.005]
- [4] Chung SY, Kim JS, Kim M, Hong MK, Lee JO, Kim CM, et al. Survey of nitrate and nitrite contents of vegetables grown in Korea. *Food Addit Contam.* 2003; 20(7):621-8. [DOI:10.1080/0265203031000124146] [PMID]
- [5] Taghipour H, Hemmati S, Faramarzi E, Somi MH, Dastgiri S, Nowrouze P. Determination of nitrate concentration in consumed vegetables and estimation of that's dietary intake in Shabestar and Khameneh City, northwest of Iran: Azar Cohort study. *Progress Nutr.* 2019; 21(S 1):336-40. <https://azarcohort.tbzmed.ac.ir/document.pdf>
- [6] WHO. Evaluation of certain food additive and contaminants. World Health Organ Tech Rep Ser. 2011; (960):1-226. [PMID]
- [7] WHO. Evaluation of certain food additives and contaminants: Forty-fourth report of the joint FAO/WHO expert committee on food additives. Geneva: WHO; 1995. <https://apps.who.int/iris/handle/10665/37246>

- [8] Hassani Moghaddam E, Bazdar A, Shaaban M. [Study of nitrate rate in some vegetables cultivated in Poldokhtar and Khorramabad, Lorestan province (Persian)]. *Iran J Health Environ.* 2019; 12(1):101-12. <https://www.cabdirect.org/cabdirect/abstract/20219928513>
- [9] Pirsahab M, Sharafi K, Morad M. [A survey on nitrite and nitrate levels in vegetables and cucurbits cultivated in northern and western plains of Kermanshah City in 2012 (Persian)]. *J Food Hyg.* 2013; 3(1):77-87. <https://www.sid.ir/en/Journal/ViewPaper.aspx?ID=396124>
- [10] Yeganeh M, Bazargan K, Samaee M, Ardebili MF, Tabbakhian S. [Residual nitrate in potato tuber samples collected from fields in important production areas and the main fruit and vegetables supply center of Tehran (Persian)]. *Iran J Soil Res.* 2019; 32(4):471-83. <https://www.cabdirect.org/cabdirect/abstract/20219931302>
- [11] Agency for Toxic Substances Disease Registry. Toxicological Profile for Nitrate and Nitrite. US Department of Health and Human Services, Public Health Service Atlanta, GA. 2017. <https://www.atsdr.cdc.gov/toxprofiles/tp204.pdf>
- [12] Institute of Standards and Industrial Research of Iran. Fruit, vegetable and derived products - Determination of nitrite and nitrate content - Molecular Absorption Spectrometric Method. ISIRI. 1st. Adment, 2013; No.4106.
- [13] Karimi B, Rajaie MS, Ghanadzadeh MJ, Mashayekhi M. [Performance evaluation of modified Fenton process using Nano scale zero-valent iron in nitrate reduction from aqueous solution (Persian)]. *Iran J Health Environ.* 2014; 6(4):471-80. <https://www.cabdirect.org/cabdirect/abstract/20143329821>
- [14] Dordas CA, Sioulas C. Safflower yield, chlorophyll content, photosynthesis, and water use efficiency response to nitrogen fertilization under rainfed conditions. *Ind Crops Prod.* 2008; 27(1):75-85. [DOI:10.1016/j.indcrop.2007.07.020]
- [15] Tabande L, Safarzadeh SS. [Evaluation of nitrate accumulation and factors affecting it in some leafy vegetables in Zanjan province (Persian)]. *Iran J Soil Res.* 2018; 32(2):189-201. [DOI:10.22092/IJSR.2018.117043]
- [16] Malakouti M. [Relationship between balanced fertilization and healthy agricultural products (a review) (Persian)]. *J Crop. Ecophysiol.* 2011; 4(16):133-50. <https://www.sid.ir/en/journal/ViewPaper.aspx?ID=260082>
- [17] Tajik R, Alimoradian A, Jamalian M, Shamsi M, Moradzadeh R, Ansari Asl B, et al. Lead and cadmium contaminations in fruits and vegetables, and arsenic in rice: A cross sectional study on risk assessment in Iran. *Iran J Toxicol.* 2021; 15(2):73-82. [DOI:10.32598/IJT.15.2.784.1]
- [18] Assessment E. Guidelines for exposure assessment. Federal Register. 1992; 57(104):22888-938. https://rais.ornl.gov/documents/GUIDELINES_EXPOSURE_ASSESSMENT.pdf
- [19] Baversad MS, Ghorbani H, Afyuni M, KheirAbadi H. [The potential risk assessment of heavy metals on human health in some agricultural products in Isfahan province (Persian)]. *J Water Soil sci.* 2014; 18(67):71-81. https://jstnar.iut.ac.ir/browse.php?a_id=1852&sid=1&slc_lang=en
- [20] Baghaie A, Aghili F. [Evaluation of lead and cadmium concentration of arak city soil and their non-cancer risk assessment in 2017 (Persian)]. *J Rafsanjan Univ Med Sci.* 2018; 17(8):769-80. <http://eprints.rums.ac.ir/6346/>
- [21] Samiee H, Hosseini H, Mofid V, Jannat B. [Contamination and risk assessment of total aflatoxin in Iranian rice in various food security regions (Persian)]. *Iran J Nutr Sci Food Technol.* 2020; 14(4):45-54. http://nsft.sbm.ac.ir/browse.php?a_code=A-10-3161-2&slc_lang=en&sid=1
- [22] Sadeghi E, Hashemian AH, Soltanian M, Soltanian S, Mohammadi M. [Study of nitrite and nitrate levels in meat products distributed in Kermanshah (Persian)]. *Iran Occup Health J.* 2014; 11(6):94-100. <https://www.cabdirect.org/globalhealth/abstract/20153168077>
- [23] Amarlooei A, Nazeri M, Nourmoradi H, Sayehmiri K, Khodarahmi F. [Investigation on the concentration of nitrate and nitrite in ilam ground waters (Persian)]. *J Ilam Univ Med Sci.* 2014; 22(4):34-41. <https://www.sid.ir/en/journal/ViewPaper.aspx?ID=420260>
- [24] Colla G, Kim HJ, Kyriacou MC, Roupael Y. Nitrate in fruits and vegetables. *Sci Hortic.* 2018; 237:221-38. [DOI:10.1016/j.scienta.2018.04.016]
- [25] Doomary H, Kamkar A, Sharifi H. [Study of Nitrate content and the effective factors on it in the cucumbers of Jiroft area (Persian)]. *J Vet Res.* 2017; 72(3):365-73. [DOI:10.22059/JVR.2017.202970.2446]
- [26] Ranasinghe R, Marapana R. Nitrate and nitrite content of vegetables: A review. *J Pharmacogn Phytochem.* 2018; 7(4):322-8. <https://www.phytojournal.com/archives?year=2018&vol=7&issue=4&ArticleId=4923>
- [27] León VM, Luzardo OP. Evaluation of nitrate contents in regulated and non-regulated leafy vegetables of high consumption in the Canary Islands, Spain: Risk assessment. *Food Chem Toxicol.* 2020; 146:111812. [DOI:10.1016/j.fct.2020.111812] [PMID]
- [28] Tamme T, Reinik M, Roasto M, Juhkam K, Tenno T, Kiis A. Nitrates and nitrites in vegetables and vegetable-based products and their intakes by the Estonian population. *Food Addit Contam.* 2006; 23(4):355-61. [DOI:10.1080/02652030500482363] [PMID]
- [29] Shahbazzadegan S, Hashemimajd K, Shahbazi B. [Determination of nitrate concentration of consumed vegetables and fruits in Ardabil (Persian)]. *J Ardabil Univ Med Sci.* 2010; 10(1):38-47. http://jarums.arums.ac.ir/browse.php?a_code=A-10-27-205&slc_lang=fa&sid=1
- [30] Pourmoghim M, Khoshtinat KH, Makkei AS, Fonod RK, Golestan B, Pirali M. [Determination of nitrate contents of lettuce, tomatoes and potatoes on sale in Tehran central fruit and vegetable market by HPLC (Persian)]. *Iran J Nutr Sci Food Technol.* 2010; 5(1):63-70. <https://www.cabdirect.org/cabdirect/abstract/20103280822>
- [31] Alemzadeh Ansari N, Sharif Manesh D, Mousavi fard S. [Evaluation of the accumulation of nitrate and nitrite in vegetables grown in different regions of Ahvaz (Persian)]. *J Plant Prod.* 2019; 42(3):333-44. [DOI:10.22055/PPD.2019.26457.1626]
- [32] Bose B, Srivastava H. Absorption and accumulation of nitrate in plants: Influence of environmental factors. *Indian J Exp Biol.* 2001; 39(2):101-10. [PMID]
- [33] Habben J. Quality constituents of carrots *Daucus Carota L.* as influenced by nitrogen and potassium fertilization. Symposium on Nutrition and Fertilization of Vegetables 29. 1971. [DOI:10.17660/ActaHortic.1973.29.22]
- [34] Pavlou GC, Ehaliotis CD, Kavvadias VA. Effect of organic and inorganic fertilizers applied during successive crop seasons on growth and nitrate accumulation in lettuce. *Sci Hortic.* 2007; 111(4):319-25. [DOI:10.1016/j.scienta.2006.11.003]

- [35] van Maanen JM, van Dijk A, Mulder K, de Baets MH, Menheere PC, van der Heide D, et al. Consumption of drinking water with high nitrate levels causes hypertrophy of the thyroid. *Toxicol Lett.* 1994; 72(1-3):365-74. [DOI:10.1016/0378-4274(94)90050-7]
- [36] McColl K. When saliva meets acid: Chemical warfare at the oesophagogastric junction. *Gut.* 2005; 54(1):1-3. [DOI:10.1136/gut.2004.047126] [PMID] [PMCID]
- [37] Eichholzer M, Gutzwiller F. Dietary nitrates, nitrites, and N-nitroso compounds and cancer risk: A review of the epidemiologic evidence. *Nutr Rev.* 1998; 56(4 Pt 1):95-105. [DOI:10.1111/j.1753-4887.1998.tb01721.x] [PMID]
- [38] Ministry of Health and Medical Education. Optimal food basket for Iranian community. In: Improvement of Nutritional Center. Publisher: Ministry of Health and Medical Education; 2013.
- [39] Uddin R, Thakur MU, Uddin MZ, Islam GR. Study of nitrate levels in fruits and vegetables to assess the potential health risks in Bangladesh. *Sci Rep.* 2021; 11(1):4704. [DOI:10.1038/s41598-021-84032-z] [PMID] [PMCID]
- [40] Mehri F, Heshmati A, Moradi M, Khaneghah AM. The concentration and health risk assessment of nitrate in vegetables and fruits samples of Iran. *Toxin Rev.* 2019. [DOI:10.1080/15569543.2019.1673424]
- [41] Thomson B, Nokes C, Cressey P. Intake and risk assessment of nitrate and nitrite from New Zealand foods and drinking water. *Food Addit Contam.* 2007; 24(2):113-21. [DOI:10.1080/02652030600934206] [PMID]
- [42] WHO. Environmental Health Criteria. Nitrites and N-nitroso compounds. Geneva: International Programme on Chemical Safety. 1978; 1-98.